

Why athletes are more likely to need pacemakers in old age

May 13 2014

A new study by The University of Manchester has shed light on why athletes are more likely to have abnormal heart rhythms.

Elderly <u>athletes</u> with a lifelong history of training and competing in endurance events like marathons, triathlons and iron man challenges can have heart rhythm disturbances, known as arrhythmias.

The Manchester research in rodents, funded by the British Heart Foundation, shows <u>molecular changes</u> in the heart's pacemaker occur in response to exercise training.

The finding, reported in *Nature Communications*, overturns the commonly held belief that an increased activity of the autonomic nervous system causes this specific reaction to endurance training.

While normal adults have resting <u>heart rates</u> between 60-100 beats per minute, hearts of endurance athletes can beat only 30 times per minute or even lower at night time when there can be long pauses between heart beats.

Cyclists Sir Chris Hoy and Miguel Indurain reportedly had resting heart rates of 30 and 28 beats per minute.

Dr Alicia D'Souza, from The University of Manchester and first author on the paper, said: "The heart rate is set by the heart's pacemaker, but this is controlled by the nervous system. The 'vagal' nerves lower the



heart rate and therefore it was assumed the low heart rate of athletes is the result of over activity of the vagal nerves.

"But our research shows this is not the case. Actually the heart's pacemaker changes in response to training and in particular there is a decrease in an important pacemaker protein, known as HCN4, and this is responsible for the low heart rate."

The researchers say these molecular changes in the sinus node - the cardiac structure responsible for generating heart rhythm - may help us to understand the more frequent occurrence of <u>heart rhythm</u> <u>disturbances</u> or even loss of consciousness in athletes.

Professor Mark Boyett, lead researcher on the study, added: "This is important because although normally a low resting heart rate of an athlete does not cause problems, elderly athletes with a lifelong training history are more likely to need an artificial electronic pacemaker fitted."

More than 500 marathons take place in Europe and America each year with around one million participants. The number of people taking part is set to increase by 5% each year.

But Professor Boyett said: "Although endurance exercise training can have harmful effects on the heart, it is more than outweighed by the beneficial effects."

Professor Jeremy Pearson, Associate Medical Director at the British Heart Foundation, said: "This study shows the heart's electrical wiring changes in mice that exercise for long periods, and these changes in heart rhythm are sustained afterwards.

"If the findings are reproduced in humans they could have implications for <u>heart</u> health in older athletes. But much more research is needed



before we could draw that conclusion."

More information: Paper: <u>dx.doi.org/10.1038/ncomms4775</u>

Provided by University of Manchester

Citation: Why athletes are more likely to need pacemakers in old age (2014, May 13) retrieved 8 May 2024 from <u>https://medicalxpress.com/news/2014-05-athletes-pacemakers-age.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.