


Risk of serious complications in obstetric anesthesia is low

May 29 2014



The risk of serious complications in obstetric anesthesia is low, according to research published in the June issue of *Anesthesiology*.

(HealthDay)—The risk of serious complications in obstetric anesthesia is low, according to research published in the June issue of *Anesthesiology*.

Robert D'Angelo, M.D., of the Wake Forest University School of Medicine in Winston-Salem, N.C., and colleagues analyzed data from nearly 257,000 patients who received neuraxial or [general anesthesia](#) for delivery. The authors sought to assess serious complications associated with obstetric anesthesia.

The researchers found that the most frequent serious complications of obstetric anesthesia were high neuraxial block, [respiratory arrest](#) in Labor and Delivery, and unrecognized spinal catheter. A serious

complication occurred in approximately one in every 3,000 episodes of obstetric anesthesia (95 percent confidence interval, one in every 2,443 episodes to one in every 3,782 episodes).

"Because serious complications related to obstetric anesthesia are rare, there were too few complications in each category to identify [risk factors](#) associated with each," the authors write. "However, since many of these [complications](#) can lead to catastrophic outcomes, it is recommended that the anesthesia provider remain vigilant and be prepared to rapidly diagnose and treat any complication."

More information: [Full Text](#)

Copyright © 2014 [HealthDay](#). All rights reserved.

Citation: Risk of serious complications in obstetric anesthesia is low (2014, May 29) retrieved 3 May 2024 from <https://medicalxpress.com/news/2014-05-complications-obstetric-anesthesia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
