

## Disease outbreak may not spur parents to have children vaccinated

May 5 2014

---

Conventional wisdom holds that when the risk of catching a disease is high, people are more likely to get vaccinated to protect themselves.

This may not be the case, however, according to a study to be presented Monday, May 5, at the Pediatric Academic Societies (PAS) annual meeting in Vancouver, British Columbia, Canada.

Researchers, led by Elizabeth R. Wolf, MD, FAAP, compared rates of infant vaccination with the diphtheria, tetanus and [pertussis](#) vaccine (DTaP) before and during an [epidemic](#) of pertussis (whooping cough) in Washington state. Surprisingly, they found no difference in [vaccination rates](#).

"We have always assumed that when the risk of catching a disease is high, people will accept a vaccine that is effective in preventing that disease. Our results may challenge this assumption," said Dr. Wolf, the Ruth L. Kirschstein National Research Service Award Fellow in General Academic Pediatrics at University of Washington, Seattle Children's Research Institute.

Washington state experienced a pertussis epidemic from Oct. 1, 2011, through Dec. 31, 2012, and infants were hit the hardest. The highly contagious bacterial disease causes uncontrollable, violent coughing that can make it hard to breathe. Pertussis also is known as whooping cough because a "whooping" sound often is heard when the patient tries to take a breath. Pertussis can lead to pneumonia, seizures (jerking and staring

spells), brain damage and death.

Dr. Wolf and her colleagues compared the proportion of 3- to 8-month-olds who had received the recommended number of doses of pertussis-containing vaccine before the epidemic and during the epidemic. Infants who received at least one dose by 3 months of age, at least two doses by 5 months and at least three doses by 7 months are considered up to date by the Advisory Committee on Immunization Practices of the Centers for Disease Control and Prevention.

"We hypothesized that a whooping cough epidemic would result in more parents getting their children immunized against whooping cough," Dr. Wolf said. "But compared to a time before the 2011-2012 whooping cough epidemic in Washington state, there was no significant increase in receipt of [whooping cough](#) vaccines for infants during the epidemic."

Results did show considerable variability in vaccination rates among different counties.

"Vaccination rates in the U.S. are still below public health goals," Dr. Wolf noted. "We don't fully understand what improves vaccine acceptance. This study found no significant increase in vaccination coverage statewide during the 2011-2012 pertussis epidemic. This finding may challenge the assumption that vaccine acceptance uniformly increases when risk of disease is high."

**More information:** Dr. Wolf will present "The Effect of the 2011-2012 Pertussis Epidemic on Infant DTaP Vaccination in Washington State" from 12:00-12:15 p.m. Monday, May 5. To view the study abstract, go to [www.abstracts2view.com/pas/view ... hp?nu=PAS14L1\\_3380.7](http://www.abstracts2view.com/pas/view...hp?nu=PAS14L1_3380.7)

Provided by American Academy of Pediatrics

Citation: Disease outbreak may not spur parents to have children vaccinated (2014, May 5)  
retrieved 18 April 2024 from

<https://medicalxpress.com/news/2014-05-disease-outbreak-spur-parents-children.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.