

Test that measures 'everyday task' performance is a good predictor of hospital readmission

May 5 2014

Patients freshly discharged from acute care hospitals with low scores on a standard test that measures how well they perform such everyday activities as moving from a bed to a chair are far more likely to need readmission to a hospital within 30 days than those who score better, according to new Johns Hopkins research.

"The Functional Independence Measure score is a direct reflection of a patient's ability to heal," says Erik Hoyer, M.D., an assistant professor in the Department of Physical Medicine and Rehabilitation at the Johns Hopkins University School of Medicine and the department's deputy director of quality and safety. "When a person cannot move his or her legs or use the bathroom independently, for example, it's telling us something about the body's physiologic reserve, its overall ability to be resilient to disease."

A report on Hoyer's research is published in the May issue of the *Journal of Hospital Medicine*.

Though many researchers have examined factors that may contribute to increased risk of rapid readmission, Hoyer says, none to his knowledge have focused on an overall standard measure of how well patients perform activities necessary for everyday living. One reason may be because few hospitals require inpatients' function to be assessed in a standardized way, but with new Medicare reimbursement guidelines in

place that cut [hospital](#) reimbursements when patients are readmitted within 30 days of discharge, that might drive interest in using similar assessments on patients in general.

Hoyer explains that all inpatient physical rehab facilities in the United States already require use of the standard Functional Independence Measure (FIM) to assess inpatients' physical capacity for a wide range of tasks, including the ability to transfer themselves between a chair and a wheelchair, the toilet or the shower; to walk, eat, bathe and dress on their own; and to effectively communicate, socially interact, solve problems and remember important information.

The FIM must be conducted within 72 hours of admission to a rehab facility, and the scores are reported to state and national agencies.

Taking advantage of this reporting requirement, Hoyer and his colleagues used various databases to pull information on 9,405 patients admitted to an inpatient rehab facility directly from an [acute care](#) hospital between July 2006 and December 2012. The data included general demographic information from each patient, such as age, gender and race/ethnicity, as well as primary diagnosis upon discharge from the hospital, the severity of illness, FIM scores upon admission to rehab and whether he or she was readmitted to an [acute care hospital](#) within 30 days of entering rehab.

Breaking the FIM scores into low, medium and high categories, the researchers found that those who scored low on the functionality test were two to three times more likely to be readmitted to the hospital within 30 days than those with high scores. Patients with low FIM scores were significantly more likely to be readmitted to the hospital even after accounting for potential confounders, including age, gender and even severity of illness.

Further investigation showed that out of the three general categories of illness that people enter rehab to treat—medical (such as congestive heart failure), neurological (such as stroke) and orthopedic (such as hip replacement)—those with medical diagnoses were the most likely to head back to the hospital within a month.

Medical diagnoses often include health problems that affect many organs at the same time, unlike neurological or orthopedic diagnoses, which affect more localized parts of the body.

"If you have a stroke or a knee replacement and can't move your legs," Hoyer explains, "it's for a very different physiological reason than if you can't move your legs because of heart failure."

Hoyer notes that getting patients more mobile and functional during their stay used to be a standard part of hospital care, and probably should be again, but because nurses and other caregivers have more and more to do in the acute care setting, that will be a challenge.

"Helping patients become more functional before they leave the hospital is important not just for Medicare policy, but also for [patients'](#) overall hospital experience and quality of life when they leave," he says. "When you talk about whether someone can use a bedpan or the toilet, it's a question of dignity."

Provided by Johns Hopkins University School of Medicine

Citation: Test that measures 'everyday task' performance is a good predictor of hospital readmission (2014, May 5) retrieved 20 April 2024 from <https://medicalxpress.com/news/2014-05-everyday-task-good-predictor-hospital.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.