

Gender may contribute to recovery time after concussion

May 6 2014

A study of concussion patients using diffusion tensor imaging (DTI) found that males took longer to recover after concussion than females did. Results of the study, which show that DTI can be used as a bias-free way to predict concussion outcome, are published online in the journal *Radiology*.

Each year, more than 17 million Americans suffer a mild traumatic brain injury (mTBI), more commonly known as a concussion, of which approximately 15 percent suffer persistent symptoms beyond three months.

Assessing outcomes and recovery time after concussion can be very subjective. Typically, physicians must rely on patient cooperation to assess injury severity.

"MRI and CT brain images of concussion patients are often normal," said Saeed Fakhraan, M.D., assistant professor of neuroradiology at the University of Pittsburgh School of Medicine. "Diffusion tensor imaging is the first imaging technique that shows abnormalities associated with concussion, because it is able to see white matter tracts at a microscopic level."

DTI is an advanced form of MRI that allows researchers to assess microscopic changes in the brain's white matter. The brain's white matter is composed of millions of nerve fibers called axons that act like communication cables connecting various regions of the brain. DTI

produces a measurement, called fractional anisotropy (FA), of the movement of water molecules along axons. In healthy white matter, the direction of water movement is fairly uniform and measures high in FA. When water movement is more random, FA values decrease. Abnormally low FA is associated with cognitive impairment in patients with brain injuries.

The research team examined the medical records and imaging results of 69 patients diagnosed with mTBI between 2006 and 2013, including 47 males and 22 females, and 21 controls consisting of 10 males and 11 females (median age of males: 17; median age of females: 16). Of the 47 males with mTBI, 32 (68 percent) were injured while playing a sport, as were 10 of the 22 females (45 percent).

All patients underwent the same evaluation, including a computerized neurocognitive test and DTI of the brain. The DTI scans of the mTBI patients revealed abnormalities within the uncinate fasciculi (UF), a [white matter](#) tract that connects the frontal and temporal lobes of the brain. Although its exact role is controversial, the UF tract is believed to allow temporal lobe-based memory associations to modify behavior through interactions with another area of the brain.

The DTI scans revealed that compared to the female mTBI patients, the male mTBI patients had significantly decreased UF FA values.

"In the future, we would like to look at the issue of gender and concussions more in depth to determine who does better and why," Dr. Fakhran said.

A statistical analysis of the data revealed that UF FA value was a stronger predictor of recovery time than initial symptom severity based on neurocognitive testing. The most substantial risk factor for a recovery time longer than three months was decreased UF FA. Male gender also

directly correlated with increased [recovery time](#).

"The potential of DTI and UF FA to predict outcome after concussion has great clinical impact," Dr. Fakhran said. "Currently, we are heavily reliant on patient reporting, and patients may have ulterior motives, such as wanting to get back to play. But you can't trick an MR scanner."

The average time to symptom recovery for all concussion patients was 54 days. However, compared to the female patients who recovered in an average of 26.3 days, recovery was significantly longer for the male patients (an average of 66.9 days), irrespective of initial symptom severity.

"Male gender and UF FA values are independent risk factors for persistent post-concussion symptoms after three months and stronger predictors of time to recovery than initial symptom severity or neurocognitive test results," Dr. Fakhran said.

He said results of the study indicate a potential role for UF FA values in triaging concussion [patients](#) in the future.

"There's prognostic value in DTI for both children participating in sports as well as for professional athletes," he said. "Lower FA values in the uncinate fasciculi could offer a metric for evaluating the severity of mild traumatic brain injuries and predicting clinical outcome. We're not at the point where DTI can provide individual prognoses yet, but that's the hope and goal."

More information: "Sex Differences in White Matter Abnormalities after Mild Traumatic Brain Injury: Localization and Correlation with Outcome," *Radiology*, 2014.

Provided by Radiological Society of North America

Citation: Gender may contribute to recovery time after concussion (2014, May 6) retrieved 27 April 2024 from

<https://medicalxpress.com/news/2014-05-gender-contribute-recovery-concussion.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.