

HRS: Bariatric surgery may prevent A-fib onset in obese

May 8 2014



(HealthDay)—Bariatric surgery may prevent the development of atrial fibrillation in morbidly obese patients, according to research presented at the annual meeting of the Heart Rhythm Society, held from May 7 to 10 in San Francisco.

Hongtao Yuan, M.D., and colleagues analyzed data for 438 patients (mean age, 59 ± 11 years; 81.1 percent female) with <u>body mass index</u> (BMI) greater than 40 kg/m^2 who received <u>bariatric surgery</u> (326 patients) or served as controls (112 patients). The authors sought to assess the effect of bariatric surgery on incidence of atrial fibrillation.

The researchers found that the group receiving bariatric surgery had a significant reduction in BMI compared with the control group ($-12.1 \pm 0.4 \text{ kg/m}^2 \text{ versus } 0.2 \pm 0.7 \text{ kg/m}^2$; P



"Bariatric surgery has a preventive effect on the incidence of <u>atrial fibrillation</u> in patients with morbid obesity," the authors write.

More information: Abstract

More Information

Copyright © 2014 HealthDay. All rights reserved.

Citation: HRS: Bariatric surgery may prevent A-fib onset in obese (2014, May 8) retrieved 5 May 2024 from https://medicalxpress.com/news/2014-05-hrs-bariatric-surgery-a-fib-onset.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.