

Study compares newer and older antipsychotic medications for schizophrenia

May 20 2014



Functional magnetic resonance imaging (fMRI) and other brain imaging technologies allow for the study of differences in brain activity in people diagnosed with schizophrenia. The image shows two levels of the brain, with areas that were more active in healthy controls than in schizophrenia patients shown in orange, during an fMRI study of working memory. Credit: Kim J, Matthews NL, Park S./PLoS One.

Among adults with schizophrenia or schizoaffective disorder, treatment with the newer, more costly antipsychotic paliperidone palmitate, compared with the older antipsychotic haloperidol decanoate, found no significant difference on a measure of effectiveness, according to a



study in the May 21 issue of JAMA.

Long-acting injectable <u>antipsychotic</u> medications are prescribed to reduce nonadherence to drug therapy and relapse in people diagnosed with a schizophrenia-spectrum disorder. The relative effectiveness of long-acting injectable versions of second-generation and older <u>antipsychotic medications</u> has not been previously assessed, according to background information in the article.

Joseph P. McEvoy, M.D., of Georgia Regents University, Augusta and colleagues randomly assigned 311 patients diagnosed with schizophrenia or schizoaffective disorder to receive monthly injections of haloperidol decanoate or paliperidone palmitate for as long as 24 months. The researchers measured rates of efficacy failure, defined as a <u>psychiatric hospitalization</u>, a need for crisis stabilization, a substantial increase in frequency of outpatient visits, a clinician's decision that oral antipsychotic could not be discontinued within 8 weeks after starting the long-acting injectable antipsychotics, or a clinician's decision to discontinue the assigned long-acting injectable due to inadequate therapeutic benefit.

The authors found no <u>significant difference</u> in the rate of efficacy failure for patients in the paliperidone palmitate group (49 [33.8 percent]) vs those in the haloperidol decanoate group (47 [32.4 percent]. The most common reasons for efficacy failure were psychiatric hospitalization and clinician discontinuation of study medication due to inadequate therapeutic effect. The researchers note that their findings do not rule out the possibility of a clinically meaningful advantage with paliperidone palmitate.

Regarding <u>adverse effects</u>, on average, participants taking paliperidone palmitate gained weight progressively over time, while those taking haloperidol decanoate lost weight. Treatment with paliperidone palmitate



was associated with greater increases in serum prolactin (a hormone), whereas haloperidol decanoate was associated with more akathisia (a movement disorder).

"The results [of this study] are consistent with previous research that has not found large differences in the effectiveness of newer and older antipsychotic medications," the authors conclude.

"Setting aside the substantial differences in cost between haloperidol decanoate and paliperidone palmitate [approximately \$35 vs. \$1,000, respectively, per injection], the results from [this] trial suggest that drug selection should be based on anticipated adverse effects rather than efficacy," writes Donald C. Goff, M.D., of the New York University School of Medicine, and Associate Editor, *JAMA*, in an accompanying editorial.

"This is true for most antipsychotics, with the notable exception of clozapine, which has established superior efficacy for treatment-resistant schizophrenia but is limited in use due to more serious adverse effects. Although patients may try medications sequentially to identify an optimal agent, this approach may be problematic if adverse effects persist after drug discontinuation, such as weight gain or involuntary movements. Additional data from patient samples exposed for a longer duration to a wider range of antipsychotics are needed to better characterize the relative risk of adverse effects, examine their longer term consequences, and identify biomarkers that will allow a personalized medicine approach. Not only is the compilation of reliable data about these drugs essential, so also is the clear communication of this information to patients as part of the shared decision-making process."

More information: <u>DOI: 10.1001/jama.2014.4310</u> <u>DOI: 10.1001/jama.2014.4311</u>



Provided by The JAMA Network Journals

Citation: Study compares newer and older antipsychotic medications for schizophrenia (2014, May 20) retrieved 2 May 2024 from https://medicalxpress.com/news/2014-05-older-antipsychotic-medications-schizophrenia.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.