

Rhythmic bursts of electrical activity from cells in ear teach brain how to hear

May 21 2014



A precise rhythm of electrical impulses transmitted from cells in the inner ear coaches the brain how to hear, according to a new study led by researchers at the University of Pittsburgh School of Medicine. They report the first evidence of this developmental process today in the online version of *Neuron*.

The ear generates spontaneous [electrical activity](#) to trigger a response in the brain before hearing actually begins, said senior investigator Karl Kandler, Ph.D., professor of otolaryngology and neurobiology, Pitt School of Medicine. These patterned bursts start at inner hair cells in the cochlea, which is part of the inner ear, and travel along the auditory nerve to the brain.

"It's long been speculated that these impulses are intended to 'wire' the brain auditory centers," he said. "Until now, however, no one has been able to provide experimental evidence to support this concept."

To map neural connectivity, Dr. Kandler's team prepared sections of a mouse brain containing the auditory pathways in a chemical that is inert until UV light hits it. Then, they pulsed laser light at a neuron, making the chemical active, which excites the [nerve cells](#) to generate an [electrical impulse](#). They then tracked the spread of the impulse to adjacent cells, allowing them to map the network a neuron at a time.

All mice are born unable to hear, a sense that develops around two weeks after birth. But even before hearing starts, the ear produces rhythmic bursts of electrical activity which causes a broad reaction in the brain's auditory processing centers. As the beat goes on, the brain organizes itself, pruning unneeded connections and strengthening others. To investigate whether the beat is indeed important for this reorganization, the team used genetically engineered mice that lack a key receptor on the inner hair cells which causes them to change their beat.

"In normal mice, the wiring diagram of the brain gets sharper and more efficient over time and they begin to hear," Dr. Kandler said. "But this doesn't happen when the [inner ear](#) beats in a different rhythm, which means the [brain](#) isn't getting the instructions it needs to wire itself correctly. We have evidence that these mice can detect sound, but they have problems perceiving the pitch of sounds."

In humans, such subtle hearing deficits are associated with Central Auditory-Processing Disorders (CAPD), difficulty processing the meaning of sound. About 2 to 3 percent of children are affected with CAPD and these children often have speech and language disorders or delays, and learning disabilities such as dyslexia. In contrast to causes of hearing impairments due to ear deficits, the causes underlying CAPD

have remained obscure.

"Our findings suggest that an abnormal rhythm of electrical impulses early in life may be an important contributing factor in the development of CAPD. More research is needed to find out whether this also holds true for humans, but our results point to a new direction that is worth following up," Dr. Kandler said.

Provided by University of Pittsburgh Schools of the Health Sciences

Citation: Rhythmic bursts of electrical activity from cells in ear teach brain how to hear (2014, May 21) retrieved 19 April 2024 from <https://medicalxpress.com/news/2014-05-rhythmic-electrical-cells-ear-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.