

Can vitamin D slow heart complications from diabetes?

May 12 2014, by Jim Dryden



Carlos Bernal-Mizrachi, MD, works with diabetes patient Helen Randall to see whether treatment with vitamin D may help lower her risk for coronary heart disease.

(Medical Xpress)—Researchers at Washington University School of Medicine in St. Louis are evaluating whether vitamin D can slow the development of cardiovascular problems in African-Americans with diabetes. They are more likely to die from cardiovascular disease than

Caucasians with diabetes.

The researchers are seeking to enroll about 90 African-Americans who are 45 to 80 years old and have type 2 diabetes. Study volunteers must not have heart disease or have suffered a stroke.

"Cardiovascular disease is a major health problem and cause of mortality in African-Americans," said Carlos Bernal-Mizrachi, MD, the study's principal investigator. "Compared to Caucasians, African-Americans suffer disproportionately from type 2 diabetes and heart disease."

Nationally, African-Americans with diabetes are 36 percent more likely than Caucasians with diabetes to die of cardiovascular disease. In past studies, Bernal-Mizrachi has found that low levels of vitamin D can double the risk of [cardiovascular disease](#) in patients with [type 2 diabetes](#). His research also shows that blood vessels near the heart are less likely to clog in people who get adequate levels of the key vitamin.

Study volunteers will be screened at the School of Medicine to determine their vitamin D status. Those with low vitamin D levels will be asked to return for a second assessment, during which investigators will evaluate participants' risk for [coronary heart disease](#) by measuring electrolyte levels, kidney function, blood cell counts, average [blood sugar levels](#) and cholesterol levels.

"In general, vitamin D deficiency is more common among African-Americans, and that may help explain why they are so much more likely than Caucasians to have severe cardiovascular complications related to diabetes," said Bernal-Mizrachi, associate professor of medicine and of cell biology and physiology.

Volunteers who meet the study's criteria will be randomly assigned to one of two groups. Those in the first group will take 600 international

units of vitamin D daily to try to raise their levels of the vitamin to the standard recommended for good health. Participants in the second study group will take 4,000 international units of the vitamin daily.

Subjects will be evaluated about every two months to measure blood sugar levels, hypertension, heart function and various markers of inflammation—all risk factors for [heart disease](#).

"Compared to other potential interventions, vitamin D is very inexpensive and doesn't have many known side effects," said Bernal-Mizrachi. "We believe vitamin D could have a very big impact if the study shows that it can prevent or delay heart problems in African-Americans with [diabetes](#)."

Provided by Washington University School of Medicine in St. Louis

Citation: Can vitamin D slow heart complications from diabetes? (2014, May 12) retrieved 24 April 2024 from

<https://medicalxpress.com/news/2014-05-vitamin-d-heart-complications-diabetes.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--