

Researchers analyze AUDs, sexual behavior among South African men

June 21 2014

In a study of South African men who drink alcohol in informal drinking environments or "shebeens," researchers from The Miriam Hospital have found a high prevalence of Alcohol Use Disorder (AUD) that directly correlates to unprotected sex. Findings support the need for interventions targeting both alcohol and HIV-risk behaviors among South African men who drink alcohol in alcohol-serving venues. The study and its findings are published in the July issue of *Drug and Alcohol Dependence*.

Lori A. Scott-Sheldon, Ph.D., lead author and researcher at The Centers for Behavioral and Preventive Medicine at The Miriam Hospital, and her colleagues examined the extent to which [alcohol](#) consumption among patrons of shebeens meets criteria for AUD, a treatable, manageable chronic condition. Shebeens in South Africa are settings in which alcohol use and sexual behavior often co-occur, and the prevalence of AUDs among men frequenting public drinking venues may in part explain South Africa's high rates of HIV.

"The HIV/AIDS epidemic remains a major public health concern in South Africa, home to the largest number of people living with HIV in the world," said Scott-Sheldon. "Not only do South Africans bear the heaviest HIV burden, they also have the highest levels of [alcohol consumption](#) per adult drinker worldwide."

Rates of hazardous or harmful drinking (defined as a pattern of drinking that increases the risk of adverse health events or consequences) are also high with one-third of South African adult drinkers reporting hazardous

or harmful alcohol use. Prior research in sub-Saharan Africa shows that alcohol use is associated with increased [sexual risk behaviors](#) that put people at risk for HIV. Consequently, alcohol consumption, as a contributing factor of HIV infection as well as other health consequences, is a major public health concern in South Africa.

Scott-Sheldon led a study of 763 men with an average age of 30 years old. They were recruited from townships in Cape Town, South Africa, and completed a self-administered survey that assessed alcohol use, sexual risk behaviors, and alcohol use situations and settings.

"Nearly two-thirds of men met criteria for AUD," Scott-Sheldon concluded. "Alcohol Use Disorder was associated with HIV risk such that men with AUD reported more unprotected sex than men without AUD.

"Of the men who met criteria for an alcohol use disorder," Scott-Sheldon added, "nearly one-third were classified as having a severe disorder. Men with an alcohol use disorder also reported engaging in [unprotected sex](#) more than [men](#) without a disorder."

Provided by Lifespan

Citation: Researchers analyze AUDs, sexual behavior among South African men (2014, June 21) retrieved 27 April 2024 from

<https://medicalxpress.com/news/2014-06-auds-sexual-behavior-south-african.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--