

Children exposed to secondhand tobacco, cooking smoke have high rates of pain and complications after tonsillectomy

June 1 2014

New research presented at this year's Euroanaesthesia meeting in Stockholm shows that children exposed to indoor coal-burning stoves and/or second-hand tobacco smoke are much more likely to suffer postoperative complications and excessive pain after tonsillectomies. The research is by Professor Daniel Sessler, Cleveland Clinic, Cleveland, Ohio, USA and Dr Onur Koyuncu, Mustafa Kemal University, Hatay, Turkey, and colleagues.

Almost half of the world's population uses <u>solid fuel</u> including biomass (wood, <u>crop residues</u>, and animal dung) or coal for heating and cooking. Because stoves are usually centrally positioned in homes and often the only source of heat, families tend to congregate around them and sleep nearby. Many stoves generate and release pollutants into household air including carbon monoxide. Children are also exposed to carbon monoxide via second-hand tobacco smoke, with estimates suggesting 40-70% of <u>children</u> are exposed worldwide.

Exposure to carbon monoxide results in carboxyhaemoglobin*, which may provoke <u>postoperative complications</u>. Elevated carboxyhaemoglobin concentrations may also increase pain sensitivity. The investigators therefore tested the primary hypothesis that children with high preoperative carboxyhaemoglobin concentrations have more postoperative complications and pain after tonsillectomies. Secondarily they tested the hypothesis that high carboxyhaemoglobin concentrations



are associated with more pain and painkiller use.

100 Turkish children scheduled for elective tonsillectomy were divided into low and high carbon monoxide exposure groups: carboxyhaemoglobin ≤ 3 or ≥ 4 g/dl. The primary outcome was complications during the seven days after surgery which included bronchospasm, laryngospasm, persistent coughing, desaturation (low blood oxygen), the need for re-intubation, low blood pressure, postoperative bleeding, and reoperation. Pain was evaluated with Wong-Baker Faces pain scales, and supplemental use of the painkiller tramadol was recorded for four hours postoperatively.

There were 36 patients in the low exposure group carboxyhemoglobin [mean 1.8 g/dl], and 64 patients were in the high exposure group [mean 6.4 g/dl]. Indoor coal-burning stoves were reported more often by families of the high than low carboxyhemoglobin children (89% versus 72%). Second-hand cigarette smoke exposure was reported by 54% of the families with children with high carboxyhaemoglobin, but only by 24% of the families of children with low carboxyhaemoglobin .

Complications were more common in patients with high carboxyhaemoglobin (47% versus 14% in the low group), with most occurring in the postanaesthesia care unit. Pain scores in the postanaesthesia care unit and one hour after surgery were significantly lower in the low exposure group. Similarly, tramadol use was lower in the low-exposure group at both 4 hours (3.5 vs. 6.0 mg) and 24 hours (3.5 vs. 6.0 mg).

The authors conclude: "Children exposed to substantial environmental <u>carbon monoxide</u> have more complications and more <u>pain</u> after general anaesthesia than less exposed children."



Provided by European Society of Anaesthesiology

Citation: Children exposed to secondhand tobacco, cooking smoke have high rates of pain and complications after tonsillectomy (2014, June 1) retrieved 20 May 2024 from https://medicalxpress.com/news/2014-06-children-exposed-secondhand-tobacco-cooking.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.