

# New research provides better understanding of endometriosis

June 5 2014

---

A mouse model of endometriosis has been developed that produces endometriosis lesions similar to those found in humans, according to a report published in *The American Journal of Pathology*. This model closely mirrors the human condition as an estrogen-dependent inflammatory disorder, and findings from the study suggest that macrophages present in shed endometrium contribute to the development of the lesions.

"One in 10 women of reproductive age have [endometriosis](#); it is as common as asthma or diabetes, but it can take up to seven years to diagnose and there is an unmet clinical need for better treatments with fewer side effects," reported lead investigator Erin Greaves, PhD, MRC Centre for Reproductive Health, Queen's Medical Research Institute, The University of Edinburgh, when addressing the UK Parliament regarding her research.

The lack of a readily available, low-cost, and suitable animal model has hindered progress in the field. Nonhuman primates offer a physiologically relevant model, but their use is limited by cost and ethical concerns. Rat and mouse models have the advantage of lower cost and smaller size but have several disadvantages. For example, mouse models often rely on suturing endometrial tissue onto the surface of pelvic organs since rodents do not naturally menstruate, raising the concern that tissue artificially placed in the pelvis may not simulate natural conditions or immune response.

The newly reported [mouse model](#) of endometriosis relies on the transplantation of menstrual endometrial tissue between genetically identical mice. In brief, a donor mouse is induced to undergo menstruation using estrogen and progesterone. The tissue that is shed from the uterus is removed and implanted into a recipient mouse, allowed to grow, and then removed and analyzed.

"We found that lesions recovered from a variety of sites in the peritoneum of the mice shared histologic similarities with human lesions, including the presence of hemosiderin, cytokeratin-positive epithelial cells, vimentin-positive stromal cells, and a well-developed vasculature. Most of the lesions had evidence of well-organized stromal and glandular structures," says Dr. Greaves. She noted other similarities including changes in the expression patterns of estrogen receptor  $\alpha$  and  $\beta$ , also similar to what is found in patient biopsies.

By performing experiments using mice with green fluorescent protein-labeled macrophages in reciprocal transfers with wild-type mice, the researchers obtained evidence that the macrophages present in the shed endometrium survive and create a pro-inflammatory microenvironment that contributes to the formation of endometriotic lesions. "We are excited by these findings because the contribution of macrophages present in shed endometrium to the etiology of endometriotic lesions has not been studied in previous mouse models," comments Dr. Greaves.

The researchers hope that this model will inform future studies investigating the role of immune cells and menstrual tissue on the development of endometriosis, advance the understanding of mechanisms of the disease, and allow the identification and study of novel targets for therapy.

According to The World Endometriosis Society, endometriosis affects an estimated 176 million women worldwide. It is an inflammatory

disorder where patches of endometrium-like tissue (the inner lining of the mammalian uterus) grow as [lesions](#) abnormally-located outside the uterine cavity. The tissue is thought to originate from endometrial fragments shed at menses. Characteristic inflammatory changes are seen such as increases in inflammatory mediators and tissue-resident immune cells. Women with endometriosis often complain of chronic, debilitating pelvic pain and infertility.

**More information:** "A Novel Mouse Model of Endometriosis Mimics Human Phenotype and Reveals Insights into the Inflammatory Contribution of Shed Endometrium," by Erin Greaves, Fiona L. Cousins, Alison Murray, Arantza Esnal-Zufiurre, Amelie Fassbender, Andrew W. Horne, Philippa T. K. Saunders ( [dx.doi.org/10.1016/j.ajpath.2014.03.011](https://doi.org/10.1016/j.ajpath.2014.03.011) ), published online ahead of *The American Journal of Pathology*, Volume 185/Issue 1 (July 2014)

Provided by Elsevier

Citation: New research provides better understanding of endometriosis (2014, June 5) retrieved 3 May 2024 from <https://medicalxpress.com/news/2014-06-endometriosis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---