

Is glaucoma a brain disease? Scientists find that jigsaw effect in glaucoma patients proves it is

June 6 2014



Acute angle closure glaucoma of the right eye (intraocular pressure was 42 in the right eye). Credit: James Heilman, MD/Wikipedia

Findings from a new study published in *Translational Vision Science & Technology (TVST)* show the brain, not the eye, controls the cellular process that leads to glaucoma. The results may help develop treatments for one of the world's leading causes of irreversible blindness, as well as contribute to the development of future therapies for preserving brain function in other age-related disorders like Alzheimer's.

In the *TVST* paper, *Refined Data Analysis Provides Clinical Evidence for Central Nervous System Control of Chronic Glaucomatous Neurodegeneration*, vision scientists and ophthalmologists describe how

they performed a data and symmetry analysis of 47 patients with moderate to severe glaucoma in both eyes. In glaucoma, the loss of vision in each eye appears to be haphazard. Conversely, neural damage within the brain caused by strokes or tumors produces visual field loss that is almost identical for each eye, supporting the idea that the entire degenerative process in glaucoma must occur at random in the individual eye—without brain involvement.

However, the team of investigators discovered during their analysis that as previously disabled optic nerve axons—that can lead to vision loss—recover, the remaining areas of permanent visual loss in one eye coincide with the areas that can still see in the other eye. The team found that the visual field of the two eyes fit together like a jigsaw puzzle, resulting in much better vision with both eyes open than could possibly arise by chance.

"As age and other insults to ocular health take their toll on each eye, discrete bundles of the small axons within the larger optic nerve are sacrificed so the rest of the axons can continue to carry sight information to the brain," explains author William Eric Sponsel, MD, of the University of Texas at San Antonio, Department of Biomedical Engineering. "This quiet intentional sacrifice of some wires to save the rest, when there are decreasing resources to support them all (called apoptosis), is analogous to pruning some of the limbs on a stressed fruit tree so the other branches can continue to bear healthy fruit."

According to the researchers, the cellular process used for pruning small optic nerve axons in glaucoma is "remarkably similar to the apoptotic mechanism that operates in the brains of people afflicted with Alzheimer's disease."

"The extent and statistical strength of the jigsaw effect in conserving the binocular visual field among the clinical population turned out to be

remarkably strong," said Sponsel. "The entire phenomenon appears to be under the meticulous control of the brain."

The *TVST* paper is the first evidence in humans that the brain plays a part in pruning optic nerve axon cells. In a previous study, *Failure of Axonal Transport Induces a Spatially Coincident Increase in Astrocyte BDNF Prior to Synapse Loss in a Central Target*, a mouse model suggested the possibility that following injury to the optic nerve cells in the eye, the brain controlled a pruning of those cells at its end of the nerve. This ultimately caused the injured cells to die.

"Our basic science work has demonstrated that axons undergo functional deficits in transport at central brain sites well before any structural loss of axons," said David J. Calkins, PhD, of the Vanderbilt Eye Institute and author of the previous study. "Indeed, we found no evidence of actual pruning of axon synapses until much, much later. Similarly, projection neurons in the brain persisted much longer, as well."

"This is consistent with the partial recovery of more diffuse overlapping visual field defects observed by Dr. Sponsel that helped unmask the more permanent interlocking jigsaw patterns once the eyes of his severely affected patients had been surgically stabilized," said Calkins.

Sponsel has already seen how these findings have positively affected surgically stabilized patients who were previously worried about going blind. "When shown the complementarity of their isolated right and left eye visual fields, they become far less perplexed and more reassured," he said. "It would be relatively straightforward to modify existing equipment to allow for the performance of simultaneous binocular visual fields in addition to standard right eye and left [eye](#) testing."

Authors of the *TVST* paper suggest their findings can assist in future research with cellular processes similar to the one used for pruning small

[optic nerve](#) axons in glaucoma, such as occurs in the brains of individuals affected by Alzheimer's.

"If the brain is actively trying to maintain the best binocular field, and not just producing the jigsaw effect accidentally, that would imply some neuro-protective substance is at work preventing unwanted pruning," said co-author of the *TVST* paper Ted Maddess, PhD, of the ARC Centre of Excellence in Vision Science, Australian National University. "Since [glaucoma](#) has much in common with other important neurodegenerative disorders, our research may say something generally about connections of other nerves within the [brain](#) and what controls their maintenance."

Provided by Association for Research in Vision and Ophthalmology

Citation: Is glaucoma a brain disease? Scientists find that jigsaw effect in glaucoma patients proves it is (2014, June 6) retrieved 19 April 2024 from <https://medicalxpress.com/news/2014-06-glaucoma-brain-disease-scientists-jigsaw.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.