

Single tick bite can pack double pathogen punch

June 20 2014



This is a female adult blacklegged tick on a hand. Credit: Kelly Oggenfuss

People who get bitten by a blacklegged tick have a higher-than-expected chance of being exposed to more than one pathogen at the same time.

The new research, published online today in the journal *PLOS ONE*, was conducted by scientists at Bard College, Sarah Lawrence College, and

the Cary Institute of Ecosystem Studies.

"We found that ticks are almost twice as likely to be infected with two [pathogens](#)—the bacterium that causes Lyme disease and the protozoan that causes babesiosis—than we would have expected," said Felicia Keesing, a professor of biology at Bard College, Adjunct Scientist at the Cary Institute, and co-author of the paper. "That means [health care providers](#) and the public need to be particularly alert to the possibility of multiple infections coming from the same tick bite."

Almost 30 percent of the ticks were infected with the agent of Lyme disease. One-third of these were also infected with at least one other pathogen. The agents of Lyme disease and babesiosis were found together in 7 percent of ticks.

The researchers collected thousands of [blacklegged ticks](#) from over 150 sites in Dutchess County, New York, an area with high incidence of tick-borne illnesses. They also collected ticks that had fed on different kinds of wildlife, including birds, rodents, opossums, and raccoons. Ticks acquire pathogens from feeding on infected hosts. DNA from each tick was extracted and tested for the presence of several pathogens.



This is a blacklegged tick under a microscope in the Ostfeld Lab. Credit: Ostfeld Lab

"Mice and chipmunks are critical reservoirs for these two pathogens, so ticks that have fed on these animals are much more likely to be co-infected," said Michelle Hersh, an assistant professor of biology at Sarah Lawrence College, past postdoctoral researcher at the Cary Institute, and lead author of the study.

"Mice and other small mammals are often particularly abundant in habitats that have been fragmented or degraded by human activity," said Richard Ostfeld of the Cary Institute of Ecosystem Studies. "That means these patterns of co-infection might get worse through time as humans continue to impact forest ecosystems."

The researchers also considered another emerging pathogen, *Anaplasma phagocytophilum*, the bacterium that causes anaplasmosis in humans. Fortunately, ticks were not more likely than expected to be co-infected with *Anaplasma* and the Lyme disease bacterium.

The researchers determined the proportion of [ticks](#) infected with each pathogen individually, then calculated the rates of co-infection expected by chance alone. Not only was co-infection with the agents of Lyme disease and babesiosis greater than expected, but rates of triple infection with the agents of Lyme, babesiosis, and anaplasmosis were about twice as likely as expected. "People in tick-infested parts of the United States such as the Northeast, Mid-Atlantic, and Upper Midwest, are vulnerable to being exposed to two or three diseases from a single tick bite," said Keesing. "And, of course, that risk increases when they're bitten by more than one tick."

More information: [dx.plos.org/10.1371/journal.pone.0099348](https://doi.org/10.1371/journal.pone.0099348)

Provided by Cary Institute of Ecosystem Studies

Citation: Single tick bite can pack double pathogen punch (2014, June 20) retrieved 25 April 2024 from <https://medicalxpress.com/news/2014-06-pathogen.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--