

Treat patients with addiction during, after hospitalization, researchers say

June 30 2014

The results of a new study demonstrate that starting hospitalized patients who have an opioid (heroin) addiction on buprenorphine treatment in the hospital and seamlessly connecting them with an outpatient office based treatment program can greatly reduce whether they relapse after they are discharged.

Led by researchers at Boston Medical Center (BMC), the study shows the important role that providers play in offering these patients [addiction treatment](#) both while in the [hospital](#) and after – even if their primary reason for being in the hospital is not for their addiction.

In this study, 139 hospitalized individuals with opioid addiction who were not already in treatment were randomized into two groups. One group received a tapered dose treatment of buprenorphine for withdrawal and referral information about community treatment programs and the other were initiated on buprenorphine, an opioid substitute proven to treat opioid addiction, along with referral to a primary care office-based buprenorphine treatment program.

Buprenorphine, which was approved by the Food and Drug Administration in 2002 for the treatment of opioid addiction, is taken orally and helps to curb opioid withdrawal symptoms.

Of those in the [buprenorphine](#) maintenance group, more than one third (37 percent) reported no illicit opioid/drug use for the month after they left the hospital compared to less than one in ten (nine percent) among the control group. These patients also reported, on average, fewer days

of illicit drug use and continued to use less over the following six months. This effect was evident despite the fact that these patients did not initially come to the hospital seeking treatment for their addiction.

"Unfortunately, referral to substance abuse [treatment](#) after discharge is often a secondary concern of physicians caring for hospitalized [patients](#)," said Jane Liebschutz, MD, MPH, a physician in general internal medicine at BMC and associate professor of medicine at Boston University School of Medicine, who served as the study's corresponding author. "However, our results show that we can have a marked impact on patient's [addiction](#) by addressing it during their hospitalization."

More information: This study is published in *JAMA Internal Medicine*.

Provided by Boston University Medical Center

Citation: Treat patients with addiction during, after hospitalization, researchers say (2014, June 30) retrieved 26 April 2024 from <https://medicalxpress.com/news/2014-06-patients-addiction-hospitalization.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
