

Study reverses current thought on treatment of cirrhosis

June 19 2014

Researchers at Mayo Clinic released a new study reversing current thought on the treatment of cirrhotic patients with type 2 diabetes. The study found that the continuation of metformin after a cirrhosis diagnosis improved survival rates among diabetes patients. Metformin is usually discontinued once cirrhosis is diagnosed because of concerns about an increased risk of adverse effects associated with this treatment in patients with liver impairment. The Mayo Clinic study was recently published in *Hepatology*.

Cirrhosis is scarring of the liver caused by forms of liver diseases, such as chronic viral hepatitis, chronic alcohol abuse and non-alcoholic fatty liver disease. This condition is the consequence of damage done to the liver over many years. As cirrhosis progresses, more and more scar tissue forms, impeding proper liver functions.

Metformin is used to treat high blood sugar levels caused by [type 2 diabetes](#). This type of diabetes works in two ways. First, it inhibits the pancreas from producing sufficient insulin, which normally regulates the movement of sugar, glucose, into cells. Glucose is the body's main source of fuel. Second, in type 2 diabetes the liver, muscle, and fat tissues become more resistant to the effects of insulin. The combination of decreased insulin production and insulin resistance results in an abnormally high level of glucose in the blood.

Obesity-related [fatty liver disease](#) can lead to liver inflammation and cirrhosis, and also is associated with diabetes. Therefore, type 2 diabetes

is found in 37 percent of cirrhotic [patients](#), five times more than in those without cirrhosis.

In the study, a sample of 172 patients continued taking [metformin](#), while another 78 individuals discontinued metformin after cirrhosis diagnosis. Patients who continued metformin as part of their treatment had a significantly longer median survival than those who stopped taking the drug. During the follow up, it was discovered that none of the patients taking metformin developed lactic acidosis, which was thought to be a common side effect of the drug in patients with cirrhosis.

"Our study suggests that metformin can be used safely in [cirrhotic patients](#). Diabetic patients who take metformin to control their [blood sugar levels](#) can continue taking metformin after cirrhosis diagnosis, if there is no specific contradiction," says gastroenterologist Lewis Roberts, M.B., Ch.B., Ph.D., senior author of the study.

With the potential implications for a major change in current clinical practice, the researchers plan to collaborate with more institutions and use nationwide databases to further validate the beneficial effects of metformin, according to Xiaodan Zhang, D.D.S., co-author of the study.

More information: "Continuation of metformin use after a diagnosis of cirrhosis significantly improved survival of patients with diabetes." Xiaodan Zhang, William S. Harmsen, Teresa A. Mettler, W. Ray Kim, Rosebud O. Roberts, Terry M. Therneau, Lewis R. Roberts and Roongruedee Chaiteerakij. *Hepatology*. 2014 May 3. [DOI: 10.1002/hep.27199](#)

Provided by Mayo Clinic

Citation: Study reverses current thought on treatment of cirrhosis (2014, June 19) retrieved 6 May 2024 from <https://medicalxpress.com/news/2014-06-reverses-current-thought-treatment-cirrhosis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.