


Over half of seniors plagued by incontinence, CDC says

June 25 2014, by Steven Reinberg, Healthday Reporter



Older men and women prone to urinary and bladder problems, report finds.

(HealthDay)—More than 50 percent of older Americans struggle with incontinence, a new government report released Wednesday shows.

"We found that half the population experienced urinary leakage or accidental bowel leakage, and about 25 percent had moderate, severe or very severe urinary leakage. And about 8 percent had moderate, severe or very severe bowel leakage," said lead researcher Yelena Gorina, a statistician at the U.S. Centers for Disease Control and Prevention's National Center for Health Statistics.

One expert noted that the impact of incontinence is significant.

"Bladder and bowel incontinence is a highly prevalent disease that has emotional, [health](#), social and economic impacts in the daily life of our

elderly population in the U.S.," said Dr. Farzeen Firoozi, a urologist at North Shore-LIJ Health System in Manhasset, N.Y.

Incontinence occurs when muscles are too weak or too active. If the muscles are weak, patients may have accidents. If muscles become too active, there may be a strong urge to go to the bathroom. There are other causes of incontinence, such as prostate problems and nerve damage.

Treatment depends on the type of problem and may include simple exercises, medicines, special devices or procedures or surgery.

Dr. Tomas Griebeling, a professor of urology at the University of Kansas and a spokesman for the American Urology Association, said incontinence increases with age.

However, "incontinence should really not be considered either a normal or inevitable part of aging," he added.

Griebeling noted that although the percentage of adults with incontinence has remained the same, as the population ages there will be more people with the problem.

According to the report, nearly 51 percent of people aged 65 and older living at home reported bladder and/or bowel incontinence. Bladder incontinence was reported by just under 44 percent and [bowel incontinence](#) by just over 17 percent.

About 50 percent of women and 25 percent of men said they suffered from bladder incontinence. White women were almost twice as likely to have bladder incontinence compared with black women, the researchers noted.

For people getting [home health care](#), there was no significant difference

in rates of incontinence by age, race and education. No matter where a patient lived, 45 percent of those getting [home health](#) care reported having difficulty with bladder and/or bowel control.

Women getting home [health care](#) were 1.7 times more likely to have bladder incontinence compared with men, the investigators found.

The CDC notes that there is a significant cost linked with incontinence. For example, in 2000 bladder incontinence cost adults an estimated \$19.5 billion, and in 2010, the average cost for bowel [incontinence](#) was estimated at \$4,100 per person.

The National Center for Health Statistics report was published in the June edition of the CDC's *Vital and Health Statistics*.

More information: Visit the [U.S. National Library of Medicine](#) for more on incontinence.

Copyright © 2014 [HealthDay](#). All rights reserved.

Citation: Over half of seniors plagued by incontinence, CDC says (2014, June 25) retrieved 26 April 2024 from

<https://medicalxpress.com/news/2014-06-seniors-plagued-incontinence-cdc.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--