

# Survey sheds light on common clinical practice for incompletely resected lung cancer

June 10 2014

---

A landmark survey of more than 700 specialists provides crucial real-world insight into the treatments most oncologists choose for lung cancer patients whose tumour has been incompletely resected, an expert from the European Society for Medical Oncology (ESMO) says.

Jean Yves Douillard, from the ICO Institut de Cancerologie de l'Ouest René Gauducheau, France, Chair of the ESMO Educational Committee, was commenting on a paper published in the journal Lung Cancer. In the study, researchers led by Raffaele Califano of The Christie NHS Foundation Trust, Manchester, UK, surveyed 768 [oncologists](#) from 41 European countries about the treatments they offered patients who had "R1 resected" non-small-cell [lung cancer](#).

R1 resection is a term used by oncologists to indicate that it is possible to find microscopic evidence of cancer cells remaining after a cancer has been surgically removed.

"We know that incomplete resection, or R1 resection, is associated with a higher risk of relapse but there are currently no strong evidence-based recommendations on how to treat these patients after surgery," Douillard says.

"This study is important since it provides a good overview on how the problem is handled in clinical practice all over Europe by practitioners who treat lung cancer."

Overall, 83% of experts surveyed were medical oncologists —specialists trained to treat cancer using chemotherapy, targeted therapies, immunotherapy and other medications.

Of the respondents, 91.4% prescribed chemotherapy, mostly cisplatin/vinorelbine or cisplatin/gemcitabine. The survey showed that the majority of doctors (85%) discussed with the patient the fact that there was limited clinical evidence to guide treatment options. Almost 50% of participants prescribed radiotherapy, with radiation oncologists most likely to offer this treatment approach.

"Treating physicians clearly believe in what they do, and try to provide the best for their patients," says Douillard. "According to the survey, however, practice is heterogeneous and varies according to the specialty of the treating physician—whether they are radiation oncologists or medical oncologists. This is why treatment decisions are best made by multidisciplinary teams."

The evidence gathered in this survey is supported by the recommendations of the 2nd ESMO Consensus Conference on Lung Cancer held in 2013, Douillard notes. That group of worldwide recognised experts recommended adjuvant chemotherapy and adjuvant radiation in R1 resected patients.

The authors of the latest paper call for prospective trials to be undertaken to provide stronger evidence to guide post-surgery treatment in this situation. Douillard agrees that such trials would be informative.

"However, trials of adjuvant treatment in R1 resected lung cancer would be very difficult to design and perform, as this is fortunately an infrequent occurrence. R1 resection would also need to be clearly defined in such studies, as it actually represents a quite heterogeneous group."

"Based on evidence from clinical trials in resected patients in whom all tumour cells have been completely removed, there is a rationale for using both chemotherapy and radiotherapy in R1-resected [non-small-cell lung cancer](#)," Douillard says.

"As the authors of this survey state, definitive proof would come from a randomised clinical trial, although such studies would be difficult to perform."

**More information:** "Use of adjuvant chemotherapy (CT) and radiotherapy (RT) in incompletely resected (R1) early stage Non-Small Cell Lung Cancer (NSCLC): A European survey conducted by the European Society for Medical Oncology (ESMO) Young Oncologists Committee." R. Califano, M.V. Karamouzis, S. Banerjee, E. de Azambuja, V. Guarneri, M. Hutka, K. Jordan, K. Kamposioras, E. Martinelli, J. Corral, S. Postel-Vinay, M. Preusser, L. Porcu, V. Torri. *Lung Cancer* - July 2014 (Vol. 85, Issue 1, Pages 74-80, [DOI: 10.1016/j.lungcan.2014.03.010](#))

Provided by European Society for Medical Oncology

Citation: Survey sheds light on common clinical practice for incompletely resected lung cancer (2014, June 10) retrieved 11 May 2024 from <https://medicalxpress.com/news/2014-06-survey-common-clinical-incompletely-resected.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.