

New anesthesia technique helps show cause of obstruction in sleep apnea

July 29 2014

A simplified anesthesia procedure may enable more widespread use of preoperative testing to demonstrate the cause of airway obstruction in patients with severe sleep apnea, suggests a study in *Anesthesia & Analgesia*.

Dr. Joshua H. Atkins and Dr. Jeff E. Mandel of the University of Pennsylvania and their colleagues have developed a new "ramp control" [anesthetic](#) technique for putting patients to sleep briefly—just enough to show the "obstructive anatomy" responsible for [sleep apnea](#). The simplified technique requires no special expertise and limits drops in [blood oxygen](#) level during testing.

Simplified Approach to Testing Before Apnea Surgery

The researchers evaluated their anesthetic technique in 97 patients participating in a study of robot-assisted surgery for severe sleep apnea. Visualizing the site of the obstruction in the upper airway is an important part of planning for apnea surgery. This can be done using a procedure called "drug-induced sleep endoscopy" (DISE), in which stepwise doses of anesthetic doses are given to put the patient to sleep and reproduce the [airway obstruction](#) causing apnea.

However, it's challenging to achieve just the right anesthetic dosage—enough to cause typical sleep-related obstruction without

causing prolonged unconsciousness or causing a deep drop in blood oxygen level (oxygen saturation). The standard technique for DISE is time-consuming and not well-suited for widespread clinical use.

In the new ramp control approach, a computerized algorithm is used to calculate the two-dose sequence of anesthetic administration likely to produce sedation in each individual patient. By contrast, the standard stepwise approach to DISE uses a sequence of up to nine doses.

The 97 patients studied had severe sleep apnea, with a median of 48 apnea-hypopnea events (complete or partial interruptions of breathing) per hour on standard testing in the sleep laboratory. The patients were being evaluated for surgery after receiving no benefit from continuous positive airway pressure (CPAP), the standard treatment for sleep apnea.

The ramp-control DISE approach allowed doctors to see and photograph the obstructive anatomy in all 97 patients. The median time to put the patient to sleep and demonstrate the cause of obstruction was just under four minutes.

Just as important, the necessary level of sedation was achieved without an undue drop in oxygen saturation levels. The median lowest oxygen saturation level during DISE was about 91 percent, compared to an 81 percent value recorded in standard sleep studies. (Normal oxygen saturation is between 95 and 100 percent.)

Two-Dose Technique May Promote Wider Use of DISE

Although it's not new, DISE is still considered a "niche procedure"—not commonly performed by most anesthesiologists. It's easy to "overshoot" the anesthetic dose; this may lead to drops in oxygen saturation,

requiring airway support steps to maintain the patient's blood oxygen level. Dr Atkins and colleagues note that their study included a "challenging" population of obese patients with severe sleep apnea, who have often been excluded from studies of other anesthetic techniques for sleep apnea.

The ramp-control DISE technique is efficient, administering just enough anesthetic to produce the desired level of sedation within a few minutes; and safe, with relatively small drops in blood [oxygen saturation](#) level. "The effective sedation seen in this study with a low rate of desaturation and infrequent need for airway support is an important result," Dr Atkins and coauthors write.

By simplifying the steps to anesthesia administration, the researchers believe their simplified approach will help to make DISE more widely available for evaluation of patients being considered for sleep apnea surgery. However, more research will be needed to determine how well the new technique can be generalized to everyday clinical practice at busy surgical centers.

More information: *Anesthesia & Analgesia*. [DOI: 10.1213/ANE.0000000000000229](#)

Provided by Wolters Kluwer Health

Citation: New anesthesia technique helps show cause of obstruction in sleep apnea (2014, July 29) retrieved 23 April 2024 from <https://medicalxpress.com/news/2014-07-anesthesia-technique-obstruction-apnea.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.