

Continuous antibiotics not necessary for many children with common prenatal abnormality

July 24 2014

Up to 5 percent of all prenatal ultrasounds uncover antenatal hydronephrosis, or enlarged kidneys, the most commonly detected prenatal abnormality in the United States. Many children with this abnormality are treated continually with preventive antibiotics for the first few years of life with the hopes of preventing the condition's associated urinary tract infections. Until recently, however, little evidence existed as to the benefits of this treatment, which involves considerable cost and inconvenience for families. But a study published this week in the *Journal of Pediatric Urology* found that, in most cases, continuous antibiotics for these children are unnecessary, findings that are especially of interest amidst increasing concern regarding antibiotic overuse.

The research, conducted by clinician-scientists at Nationwide Children's Hospital and collaborating institutions, showed that the effectiveness of preventive or prophylactic [antibiotics](#) for children with a prenatal diagnosis of enlarged kidneys depends on the severity of their condition.

For parents like Katie and Drew Danielson, this news comes as a relief. The couple decided to forgo continuous preventive antibiotics for their son's mild case of vesicoureteral reflux (VUR) after consulting a pediatric urologist at Nationwide Children's. VUR is one of the primary causes of enlarged kidneys detected through prenatal ultrasound.

"Having a study like this show that it was the right thing to do is very reaffirming," Katie said. Their son Alexander is now a year old. "We didn't want our son to be on constant antibiotics for a number of reasons, and as it turns out, he wouldn't have needed them anyway. He hasn't had a single UTI."

The Danielsons' experience fits in with the study's results.

"We found that for children with ureteral dilation, severe or high-grade VUR or obstruction of the ureterovesical junction, antibiotics significantly reduced the risk of developing UTIs," said Daniel B. Herz, MD, a urologist at Nationwide Children's and lead author of the recent paper. "In those cases, prophylactic antibiotics are appropriate. But for kids with only mild problems associated with their hydronephrosis, continuous antibiotic therapy is probably not needed."

Children who did not have one of those three more serious diagnoses were unlikely to develop a UTI even without prophylactic antibiotics, Dr. Herz explained. "Because they are at low risk of infection, putting them on continuous antibiotic therapy just doesn't make sense."

The researchers compared medical records of over 400 children who were followed for at least 2 years, 278 of whom took continuous antibiotic prophylaxis and 127 of whom did not. They then tallied the number of UTIs in each group and analyzed which conditions resulted in increased UTI risk. The cases were patients from Nationwide Children's and the Children's Hospital at Dartmouth-Hitchcock.

The study comes on the heels of considerable discussion in the peer-reviewed literature regarding the proper protocol for identifying which children with such conditions could benefit from [prophylactic antibiotics](#). Recent publications are generally favoring less of a one-treatment-fits-all approach, Dr. Herz said. Instead, studies like this one support the use

of continuous antibiotics only when a child's condition truly merits it for the prevention of recurrent or serious infections. Dr. Herz also suggested that this trend may also relieve some concerns over the growing problem of antibiotic resistance due to [antibiotic overuse](#).

"Our hope is that this study will help guide physicians when they're deciding how to treat a young patient with one of these conditions," Dr. Herz said "We're getting better at identifying which kids are most at risk of UTIs and who can truly benefit from preventive continuous antibiotics. And if we can avoid the hassle and expense of continuous antibiotics for many families without causing them any harm, we should."

More information: *Journal of Pediatric Urology*, 22 July 2014. E-pub ahead of print.

Provided by Nationwide Children's Hospital

Citation: Continuous antibiotics not necessary for many children with common prenatal abnormality (2014, July 24) retrieved 19 April 2024 from <https://medicalxpress.com/news/2014-07-antibiotics-children-common-prenatal-abnormality.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
