

Controlling childbirth pain tied to lower depression risk

July 23 2014

Controlling pain during childbirth and post delivery may reduce the risk of postpartum depression, writes Katherine Wisner, M.D., a Northwestern Medicine® perinatal psychiatrist, in a July 23 editorial in *Anesthesia & Analgesia*.

Wisner's editorial is based on a new Chinese study that found women who had pain control with epidural anesthesia during a [vaginal delivery](#) had a much lower risk for [postpartum depression](#) than women who didn't have the epidural.

"Maximizing pain control in labor and delivery with your obstetrician and [anesthesia](#) team might help reduce the risk of postpartum depression," Wisner said.

The study findings are among the few to examine the relationship between pain during labor and postpartum depression.

"It's a huge omission that there has been almost nothing in postpartum depression research about pain during labor and delivery and postpartum depression," Wisner said. "There is a well-known relationship between acute and [chronic pain](#) and depression."

Wisner is director of Northwestern's Asher Center for the Study and Treatment of Depressive Disorders. She also is a professor of psychiatry and behavioral sciences and obstetrics and gynecology at Northwestern University Feinberg School of Medicine and a psychiatrist at

Northwestern Memorial Hospital.

The Chinese study found that women who had an epidural for pain relief during labor for a vaginal delivery had a 14 percent rate of depression at six weeks postpartum compared to nearly 35 percent rate of depression for those who did not have the pain relief. An epidural was the only means of pain control available to the women.

The study also found that breastfeeding was more common in the group who had an epidural for pain compared to those who did not (70 percent versus 50 percent.)

"These findings are quite exciting and further research should be done to confirm them, especially in women at increased risk of postpartum depression and in women from other cultures," Wisner said.

The incidence of severe acute postpartum pain is approximately 11 percent, Wisner reports in the article. The incidence of chronic pain varies by study but ranges from 1 to 10 percent for vaginal delivery and 6 to 18 percent after a cesarean.

Biological and emotional factors contribute to postpartum depression, which affects 14.5 percent of [women](#) who give birth. A woman who has chronic pain one to two months after delivery should be screened for depression, noted Wisner, also the Norman and Helen Asher Professor at Feinberg.

Managing acute postpartum pain supports the new mother's ability to emotionally attach and care for her infant, Wisner points out.

"Pain control gets the mother off to a good beginning rather than starting off defeated and exhausted," Wisner said. "Whether it's vaginal or cesarean section delivery, pain control postpartum is an issue for all new

mothers. There is no way to have a delivery without pain. The objective here is to avoid severe pain. Controlling that [delivery pain](#) so a woman can comfortably develop as a mother is something that makes a lot of sense."

Provided by Northwestern University

Citation: Controlling childbirth pain tied to lower depression risk (2014, July 23) retrieved 10 April 2024 from <https://medicalxpress.com/news/2014-07-childbirth-pain-tied-depression.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--