

# How you cope with stress may increase your risk for insomnia

July 3 2014

---



Credit: xiaphias/Wikipedia

A new study is the first to identify specific coping behaviors through which stress exposure leads to the development of insomnia.

Results show that coping with a stressful event through behavioral disengagement – giving up on dealing with the stress – or by using alcohol or drugs each significantly mediated the relationship between

stress exposure and [insomnia](#) development. Surprisingly, the coping technique of self-distraction – such as going to the movies or watching TV – also was a significant mediator between stress and incident insomnia. Furthermore, the study found that cognitive intrusion - recurrent thoughts about the stressor - was a significant and key mediator, accounting for 69 percent of the total effect of stress exposure on insomnia.

"Our study is among the first to show that it's not the number of stressors, but your reaction to them that determines the likelihood of experiencing insomnia," said lead author Vivek Pillai, PhD, research fellow at the Sleep Disorders & Research Center at Henry Ford Hospital in Detroit, Michigan. "While a stressful event can lead to a bad night of sleep, it's what you do in response to stress that can be the difference between a few bad nights and chronic insomnia."

Study results are published in the July 1 issue of the journal *Sleep*.

The study involved a community-based sample of 2,892 good sleepers with no lifetime history of insomnia. At baseline the participants reported the number of stressful life events that they had experienced in the past year, such as a divorce, serious illness, major financial problem, or the death of a spouse. They also reported the perceived severity and duration of each stressful event. Questionnaires also measured levels of cognitive intrusion and identified coping strategies in which participants engaged in the seven days following the [stressful event](#). A follow-up assessment after one year identified participants with insomnia disorder, which was defined as having symptoms of insomnia occurring at least three nights per week for a duration of one month or longer with associated daytime impairment or distress.

"This study is an important reminder that stressful events and other major life changes often cause insomnia," said American Academy of

Sleep Medicine President Dr. Timothy Morgenthaler. "If you are feeling overwhelmed by events in your life, talk to you doctor about strategies to reduce your stress level and improve your sleep."

According to the authors, the study identified potential targets for therapeutic interventions to improve coping responses to stress and reduce the risk of insomnia. In particular, they noted that mindfulness-based therapies have shown considerable promise in suppressing cognitive intrusion and improving sleep.

"Though we may not be able to control external events, we can reduce their burden by staying away from certain maladaptive behaviors," said Pillai.

The American Academy of Sleep Medicine reports that short-term insomnia disorder lasting less than three months occurs in 15 to 20 percent of adults and is more prevalent in women than in men.

Provided by American Academy of Sleep Medicine

Citation: How you cope with stress may increase your risk for insomnia (2014, July 3) retrieved 18 April 2024 from <https://medicalxpress.com/news/2014-07-cope-stress-insomnia.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--