

Pregnancies following egg donation associated with more than three-fold higher risk of hypertension

July 1 2014

With an ever-ageing female patient population, egg donation is an increasingly common treatment in infertility. ESHRE's own annual reports on fertility treatments in Europe show a rise in egg donation cycles from 15,028 in 2007 to 24,517 in 2010 (to 4.05% of all treatments). This proportion is still some way behind the USA, where egg donation now accounts for around 12% of all treatments.

As women age their store of viable eggs reduces such that their "ovarian reserve" (and likelihood of pregnancy) declines. Once the eggs have gone, they cannot be replaced, and egg donation is the only possible treatment. In addition, studies consistently show that eggs produced by women in older age groups form embryos with a higher prevalence of chromosome defects than those from younger women: these embryos are less likely to implant and more likely to result in miscarriage if they do implant. As a result, fertility treatment with donor eggs is much more common among older women than among younger.

Studies also show that egg donation is an increasingly successful treatment, with live birth rates of around 55% per transfer recorded in the latest US data; success largely depends on the age of the donor, not on the age of the recipient.

Egg donation, like IVF and ICSI, is considered a safe treatment, despite the older age of [patients](#), but a new study from France has found that the

pregnancies of egg donation patients are at a higher risk of disorders - particularly relating to high blood pressure - than the pregnancies of routine IVF patients using their own eggs.

Details of this study are presented today by Dr H  l  ne Letur from the Institut Mutualiste Montsouris in Paris at the 30th Annual Meeting of ESHRE held in Munich.

"A few other studies have shown results suggesting an increased risk of pregnancy-induced hypertension in egg donation patients," said Dr Letur. "However, most of them have small samples and do not adequately control for several important confounders, such women's age, multiple pregnancy, and infertility history. Our aim was to find out whether pregnancies from egg donation are genuinely associated with a higher risk of hypertension and pre-eclampsia than those from treatments using the patient's own eggs. This has growing importance because of the increasing number of egg donations."(1)

This was a case-control study from seven IVF centres in France in which each singleton egg donation pregnancy recorded at 7-8 weeks between 2005 and 2011 was matched with two control routine IVF pregnancies; 580 pregnancies were included in the study (217 egg donation, 363 controls).

Results showed that the egg donation patients had a more than three-fold higher risk of pregnancy-induced hypertension than the routine IVF patients, with prevalence rising from 5.3% to 17.8% (odds ratio 3.84, 95% CI 1.89-7.77) and an even higher risk of pre-eclampsia, from 2.8% to 11.2% (OR 4.60, 95% CI 1.81-11.67) - although the latter finding was based on very few numbers. Further analysis showed a modest effect of patient age (OR 1.08) and no significant effect of previous pregnancy, previous IVF cycles or frozen embryo transfer.(2)

"This study confirms several other reports in the literature," said Dr Letur, "with a large sample and matched control group. We would have to conclude from the results that egg donation itself is a risk factor for pregnancy-induced hypertension and pre-eclampsia."

Dr Letur added that the high prevalence of pregnancy induced hypertension in the egg donation group (almost 18% were affected) means that patients and obstetricians must be aware of the risk.

"Preventive measures and care are necessary," she said. These include screening for risk factors for hypertension (such as obesity and diabetes) and early treatment if gestational hypertension is detected. "For the future," she added, "might we match donors and recipients based on genetic factors, or consider fertility preservation for women with a high risk of premature ovarian failure?"

Possible biological explanations for the findings include changes in the immune tolerance of the embryo, whose full genome is not concordant with the recipient's. Blood pressure may rise, says Dr Letur, "to improve materno-fetal exchanges".

More information: Abstract O-163: Pregnancies issued from egg donation are associated to a higher risk of hypertensive pathologies than control ART pregnancies. Results of a large comparative cohort study

Notes

1. Pregnancy-induced hypertension (or gestational hypertension) is usually defined by a blood pressure reading of 140/90 mmHg measured on at least two occasions after 20 weeks' gestation. Pre-eclampsia is a rarer condition defined by a similar blood pressure reading plus an elevated measure of protein in a urine sample ("proteinuria"). Age (over 40, adolescence), multiple pregnancy, hypertension and family history have been identified as risk factors. Around 3-5% of all pregnancies are

affected. Most cases of pre-eclampsia are mild and cause no trouble, but deteriorating cases can be serious for both mother and baby. In this study gestational hypertension was defined as a BP reading of 140/90 mmHg without proteinuria on at least two occasions after 20 weeks' gestation; pre-eclampsia as 140/90 mmHg with proteinuria on repeated readings after 20 weeks; and eclampsia as 160/140 with proteinuria on repeated readings after 20 weeks.

2. While egg donation is generally a treatment for women of an older maternal age (58% in ESHRE's latest monitoring figures were aged over 40), the picture in France is slightly different. There, says Dr Letur, egg donation in the social security system is restricted to women under 43, and in this study the average patient age was 34.5 years. The most frequent indication was premature ovarian failure (premature menopause), suggesting that the results in this study are a genuine effect of egg donation, and not of older age.

Provided by European Society of Human Reproduction and Embryology

Citation: Pregnancies following egg donation associated with more than three-fold higher risk of hypertension (2014, July 1) retrieved 4 May 2024 from

<https://medicalxpress.com/news/2014-07-pregnancies-egg-donation-three-fold-higher.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--