


# New findings show strikingly early seeding of HIV viral reservoir

July 20 2014

---



Diagrammatical representation of the molecular structure of HIV-1 protease complexed with the inhibitor indinavir. Credit: Public Domain

The most critical barrier for curing HIV-1 infection is the presence of the viral reservoir, the cells in which the HIV virus can lie dormant for many years and avoid elimination by antiretroviral drugs. Very little has been known about when and where the viral reservoir is established

during acute HIV-1 infection, or the extent to which it is susceptible to early antiretroviral therapy (ART).

Now a research team led by investigators at Beth Israel Deaconess Medical Center (BIDMC) in collaboration with the U.S. Military HIV Research Program has demonstrated that the viral reservoir is established strikingly early after intrarectal simian immunodeficiency virus (SIV) [infection](#) of rhesus monkeys and before detectable viremia.

The findings appear online in the journal *Nature*.

"Our data show that in this animal model, the viral reservoir was seeded substantially earlier after infection than was previously recognized," explains senior author Dan H. Barouch, MD, PhD, Director of the Center for Virology and Vaccine Research at BIDMC and steering committee member of the Ragon Institute of MGH, MIT and Harvard. "We found that the reservoir was established in tissues during the first few days of infection, before the virus was even detected in the blood."

This discovery coincides with the recently reported news of the HIV resurgence in the "Mississippi baby," who was believed to have been cured by early administration of ART. "The unfortunate news of the virus rebounding in this child further emphasizes the need to understand the early and refractory viral reservoir that is established very quickly following HIV infection in humans," adds Barouch, a Professor of Medicine at Harvard Medical School.

In this new study, the scientific team initiated suppressive ART in groups of monkeys on days 3, 7, 10 and 14 after intrarectal SIV infection. Animals treated on day 3 following infection showed no evidence of virus in the blood and did not generate any SIV-specific immune responses. Nevertheless, after six months of suppressive ART, all of the animals in the study exhibited viral resurgence when treatment

was stopped.

While early initiation of ART did result in a delay in the time to viral rebound (the time it takes for [virus](#) replication to be observed in the blood following cessation of ART) as compared with later treatment, the inability to eradicate the viral reservoir with very early initiation of ART suggests that additional strategies will be needed to cure HIV infection.

"The strikingly early seeding of the viral reservoir within the first few days of infection is sobering and presents new challenges to HIV-1 eradication efforts," the authors write. "Taken together, our data suggest that extremely early initiation of ART, extended ART duration, and probably additional interventions that activate the viral reservoir will be required for HIV-1 eradication."

**More information:** Rapid seeding of the viral reservoir prior to SIV viraemia in rhesus monkeys, *Nature*, [dx.doi.org/10.1038/nature13594](https://doi.org/10.1038/nature13594)

Provided by Beth Israel Deaconess Medical Center

Citation: New findings show strikingly early seeding of HIV viral reservoir (2014, July 20) retrieved 19 April 2024 from <https://medicalxpress.com/news/2014-07-strikingly-early-seeding-hiv-viral.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--