

# Varenicline combined with nicotine patch improves smoking cessation rates

July 8 2014

---

Combining the smoking cessation medication varenicline with nicotine replacement therapy was more effective than varenicline alone at achieving tobacco abstinence at 6 months, according to a study in the July 9 issue of *JAMA*.

The combination of behavioral approaches and pharmacotherapy are of proven benefit in assisting smokers to quit. Combining [nicotine replacement therapy](#) (NRT) with varenicline has been a suggested treatment to improve smoking [abstinence](#), but its effectiveness is uncertain, according to background information in the article.

Coenraad F. N. Koegelenberg, M.D., Ph.D., of Stellenbosch University, Cape Town, South Africa, and colleagues randomly assigned 446 generally healthy smokers to nicotine or placebo patch treatment 2 weeks before a target quit date (TQD) and continued for an additional 12 weeks. Varenicline was begun 1 week prior to TQD, continued for a further 12 weeks, and tapered off during week 13. The study was conducted in 7 centers in South Africa from April 2011 to October 2012; 435 participants were included in the efficacy and safety analyses.

The researchers found that participants who received active NRT and varenicline were more likely to achieve continuous abstinence from smoking (confirmed by exhaled carbon monoxide measurements) at 12 weeks (55.4 percent vs 40.9 percent) and 24 weeks (49.0 percent vs 32.6 percent) and point prevalence abstinence (a measure of abstinence based on behavior at a particular point in time) at six months (65.1 percent vs

46.7 percent) than those receiving placebo NRT and varenicline.

In the combination treatment group, there was more nausea, sleep disturbance, skin reactions, constipation, and depression reported, with only skin reactions reaching statistical significance (14.4 percent vs 7.8 percent); the varenicline-alone group experienced more abnormal dreams and headaches.

"In this study, to our knowledge the largest study to date examining the efficacy and safety of supplementing [varenicline](#) treatment with NRT, we have found the combination treatment to be associated with a statistically significant and clinically important higher continuous abstinence rate at 12 and 24 weeks, as well as a higher point prevalence abstinence rate at 6 months," the authors write.

They add that further studies are needed to assess long-term efficacy and safety.

**More information:** [DOI: 10.1001/jama.2014.7195](https://doi.org/10.1001/jama.2014.7195)

Provided by The JAMA Network Journals

Citation: Varenicline combined with nicotine patch improves smoking cessation rates (2014, July 8) retrieved 27 April 2024 from <https://medicalxpress.com/news/2014-07-varenicline-combined-nicotine-patch-cessation.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--