


Bypass commands from the brain to legs through a computer

August 14 2014




When turning off the computer-aided spinal cord bypass, the lower extremities which were in a relaxed state did not move even if the subject was swinging his/her arms. With the bypass turned on, when the subject swung his/her arms by his/her own will and a walking motion of the lower extremities began in rhythm to the motion of the arms. Credit: © Yukio Nishimura

Gait disturbance in individuals with spinal cord injury is attributed to the interruption of neural pathways from brain to the spinal locomotor center, whereas neural circuits locate below and above the lesion maintain most of their functions. An artificial connection that bridges the lost pathway and connects brain to spinal circuits has potential to ameliorate the functional loss.

A Japanese research group led by Shusaku Sasada, research fellow and Yukio Nishimura, associate professor of the National Institute for Physiological Sciences (NIPS), National Institutes of Natural Sciences (NINS) has successfully made an artificial connection from the brain to the locomotion center in the [spinal cord](#) by bypassing with a [computer interface](#). This allowed subjects to stimulate the spinal locomotion center using volitionally-controlled muscle activity and to control walking in legs. This result was published online in *The Journal of Neuroscience* on August 13, 2014.

Neural networks in the spinal cord, locomotion center are capable of producing rhythmic movements, such as swimming and walking, even when isolated from the brain. The brain controls spinal locomotion center by sending command to the spinal locomotion center to start, stop and change waking speed. In most cases of spinal cord injury, the loss of this link from the brain to the locomotion center causes problems with walking.

The research group came up with bypassing the functioning brain and locomotion center with the computer to compensate lost pathways as a way to enable individuals with spinal cord injury to regain walking ability.



This image shows an artificial connection that connects brain to spinal circuits.
Credit: © Yukio Nishimura

Since the arm movement associate with leg movement when we walk they used muscle activity of arm to sarogate the [brain](#) activity. The computer interface allowed subjects to control magnetic stimulator that drive to the spinal locomotion center non-invassively using volitionally-controlled [muscle activity](#) and to control walking in legs. As a results of experiments in people who are neurologically intact, the subjects were asked to make own legs relaxed and passively controlled via computer

interface that was controlled by arm muscle, walking behavior in legs was induced and subjects could control the step cycle volitionally as well. However without bypassing with the computer interface, the legs did not move even if the arms muscle was volitionally activated.

"We hope that this technology would compensate for the interrupted pathways' function by sending an intentionally encoded command to the preserved spinal locomotor center and regain volitionally-controlled walking in individuals with paraplegia. However, the major challenge that this technology does not help them to dodge obstacles and to maintain posture. We are carefully working toward clinical application in near future", Nishimura said.

Provided by National Institutes of Natural Sciences

Citation: Bypass commands from the brain to legs through a computer (2014, August 14) retrieved 18 April 2024 from <https://medicalxpress.com/news/2014-08-bypass-brain-legs.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--