

Study shows childbirth complications vary widely at US hospitals

August 4 2014

Expectant mothers anticipate a smooth delivery, yet 13 percent of all women in the U.S. experience obstetrical complications. A University of Rochester-led study, published in the August issue of *Health Affairs*, shows complication rates can vary as much as five-fold among hospitals, prompting researchers to call for the development of a national quality reporting system to improve maternal outcomes for more than 4 million women who give birth each year.

Women who delivered by cesarean at low-performing hospitals experienced lacerations, hemorrhage, clots or infections at five-times the rate of high-performing hospitals—21 percent compared to 4.4 percent. Those who delivered vaginally at low-performing hospitals were twice as likely to suffer complications, 22.6 percent versus 10.4 percent at high-performing hospitals. Researchers determined the low, average or high performing hospitals based upon a calculation of the relative risk that a patient would experience a major complication.

"The key finding is that there is significant variability in maternal outcomes across US hospitals. This presents us with an opportunity to identify 'best practices' at hospitals with low rates of <u>maternal</u> <u>complications</u> in order to improve outcomes for patients in all hospitals," said lead author Laurent G. Glance, M.D., <u>health outcomes</u> researcher and vice-chair for research in Anesthesiology at the University of Rochester.

The American Congress of Obstetricians and Gynecologists (ACOG)



and the American Society of Anesthesiologists (ASA) have launched an initiative to create a platform for measuring and reporting benchmarking information on maternal outcomes. This quality reporting initiative could become a powerful tool for improving maternal outcomes in the United States.

In this study, researchers analyzed 750,000 deliveries in the 2010 Healthcare Cost and Utilization's Nationwide Inpatient Sample. However, Glance considers the findings preliminary because they are based on administrative data which lack information on potentially important risk factors. It is also important to realize that most of these complications, although important, are rarely life-threatening.

Childbirth accounts for one in four hospital discharges, resulting in \$100 billion in <u>hospital</u> charges in 2008 alone.

High-risk obstetrician J. Christopher Glantz, M.D., M.P.H., an author on the paper, said there's great value in having this information to quantify the well-known variations in practice and its impact on maternal outcomes. He is working on an American College of Obstetricians and Gynecologists initiative to reduce pregnancy-related deaths from severe hypertension, thromboembolism and hemorrhaging during pregnancy and childbirth in New York.

"In the OB field, individual practice styles, training and anecdotal experience shape how we practice, but we didn't expect to see such wide differences in maternal outcomes, which is all we studied here," said Glantz, of the University of Rochester. "For the most part babies and the mothers do well, but we can do even better by studying the hospitals that perform well and following their best practices."

Provided by University of Rochester Medical Center



Citation: Study shows childbirth complications vary widely at US hospitals (2014, August 4) retrieved 27 April 2024 from

https://medicalxpress.com/news/2014-08-childbirth-complications-vary-widely-hospitals.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.