

Eating at fast food, full service restaurants linked to more calories, poorer nutrition

August 7 2014

For adults, eating at both fast-food and full-service restaurants is associated with significant increases in the intake of calories, sugar, saturated fat, and sodium, according to a new study. The study, appearing early online in *Public Health Nutrition*, finds on days when adults ate at a restaurant, they consumed about 200 additional total daily calories whether they ate at fast- food restaurants or at full-service restaurants.

Previous studies looking at restaurant food consumption have found that adults who reported eating [fast food](#) consumed more calories, fat, and sodium, as well as fewer fruits, vegetables and vitamins compared to than those who did not report eating fast food. Studies have also linked meals consumed at both fast-food and full-service restaurants with higher caloric intake.

For the current study, Binh T. Nguyen of the American Cancer Society and Lisa M. Powell of the University of Illinois at Chicago used more recent data from more than 12,000 respondents between the ages of 20 and 64 taking part in the National Health and Nutrition Examination Survey 2003 (NHANES). Participants were asked about visits to fast-food and full-service restaurants on two successive days.

The study found on days when eating at a fast-food restaurant, there was a net increase of total [energy intake](#) (194.49 kcal), [saturated fat](#) (3.48 g), sugar (3.95 g) and sodium (296.38 mg). Eating at a full-service restaurant was also associated with an energy intake (205.21 kcal), and

with higher intake of saturated fat (2.52 g) and sodium (451.06 mg).

The study found Individual characteristics moderated the impact of restaurant food consumption. Net energy intake was larger for black adults compared with their white and Hispanic counterparts and greater for middle-income v. high-income adults.

"The United States is one of the most obese nations in the world, with more than one in three adult men and women in defined as obese," said Dr. Nguyen. "Just as obesity rates rise, there's been a marked increase in total energy consumption consumed away from home, with about one in four calories coming from fast food or full service [restaurants](#) in 2007. Our study confirms that adults' fast-food and full-service restaurant consumption was associated with higher daily total energy intake and poorer dietary indicators."

The researchers say the larger adverse effect they measured on energy intake for some lower socio-economic and minority populations has policy implications. They say efforts to improve diet and reduce energy intake from restaurant sources could actually help to reduce racial and socio-economic disparities in Americans' diets.

More information: The impact of restaurant consumption among US adults: effects on 2 energy and nutrient intakes; Public Health Nutr Published Early Online August 7, 2014; [DOI: 10.1017/S1368980014001153](#)

Provided by American Cancer Society

Citation: Eating at fast food, full service restaurants linked to more calories, poorer nutrition (2014, August 7) retrieved 3 May 2024 from <https://medicalxpress.com/news/2014-08-fast-food->

full-restaurants-linked.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.