

Food allergies more widespread among inner-city children

August 14 2014

Already known for their higher-than-usual risk of asthma and environmental allergies, young inner-city children appear to suffer disproportionately from food allergies as well, according to results of a study led by scientists at the Johns Hopkins Children's Center.

The federally funded multi-center study, described online Aug. 13 in the *Journal of Allergy and Clinical Immunology*, found that at least one in 10 [children](#) from four large U.S. cities has a food allergy. However, the true number may be even higher, the investigators say, because the study used highly stringent criteria and counted only the three most common food allergies.

"Our findings are a wake-up call, signaling an urgent need to unravel the causes, contributors and mechanisms that drive the high prevalence of food allergies among an already vulnerable group known for its high risk of asthma and environmental allergies," says senior investigator Robert Wood, M.D., director of pediatric allergy and immunology at Johns Hopkins.

Nearly 3 percent of adults and 6 percent of young children in the United States have one or more food allergies, according to the latest estimates from the National Institutes of Health. Moreover, Wood notes, food allergies among children have been on a steady rise over the last 20 years, and experts have long suspected that children in urban areas are no exception. The new study largely affirms that trend but also points to a subgroup of children who may have higher-than-average allergy risk.

For the study, the investigative team followed 516 inner-city children from birth through age 5, living in Baltimore, Boston, New York City and St. Louis. Each year of the study, the investigators measured each child's exposure to household allergens, conducted physical exams, tracked the children's diets and reviewed their health histories. The team also analyzed blood samples at 1, 2, 3 and 5 years to measure the presence of food-specific immunoglobulin E (IgE) antibodies to milk, eggs and peanuts. IgE antibodies to food proteins are immune chemicals released by the body that herald a food allergy. Based on blood test results, other clinical findings and symptoms, the investigators classified a child as allergic, possibly allergic, sensitive to a particular substance or not sensitive. Sensitization to a substance, signaled by the presence of IgE antibodies in the blood, renders a person more likely to develop allergic symptoms, but it is not sufficient to diagnose a true food allergy, which is always marked by clinical symptoms. In this study, the investigative team deemed allergic only those children who had both clinical symptoms and elevated IgE antibodies. This stringent criterion likely underestimates the true number of kids with food allergies, Wood adds.

Overall, more than half (55 percent) of children in the study were classified as sensitive to milk, eggs or peanuts. Nearly 10 percent of them met criteria for a full-blown food allergy. The most common allergy was to peanuts (6 percent), followed by eggs (4.3 percent) and milk (2.7 percent). An additional 17 percent were classified as "possibly allergic," a subgroup that had elevated IgE antibodies but no clear history of allergic reactions to peanuts, eggs or milk. Twenty-nine percent were classified as "sensitive but tolerant," a group that included those with elevated IgE antibodies and a known history of consuming allergenic foods but who were able to tolerate the foods in question without allergic symptoms.

Breastfed children appeared to have a higher risk for developing food

allergies. Children living in houses with higher levels of endotoxin, a molecule released by certain types of bacteria, were less likely to have a [food allergy](#). This latter finding, the investigators say, is consistent with the so-called hygiene hypothesis, which suggests that early-life exposure to certain microbes can play a protective role against asthma and allergies.

Children with food allergies were also more likely to suffer from [environmental allergies](#), wheezing and eczema, an allergic skin condition.

Provided by Johns Hopkins University School of Medicine

Citation: Food allergies more widespread among inner-city children (2014, August 14) retrieved 6 July 2024 from

<https://medicalxpress.com/news/2014-08-food-allergies-widespread-inner-city-children.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.