


Lipids boost the brain

August 11 2014



Top: Membranes containing monounsaturated (left) and polyunsaturated (right) lipids after adding dynamin and endophilin. In a few seconds membranes rich in polyunsaturated lipids undergo many fissions. (Credit: © Mathieu Pinot) Bottom: Transferrin (iron transport) endocytosis in cells containing polyunsaturated lipids

in their membranes (right) compared to those of cells without polyunsaturated lipids (left). In 5 minutes, the number of endocytosis vesicles formed (transferrin brought in is red) is increased by almost a factor of 10, reflecting easier endocytosis. (Credit: © H  l  ne Barelli)

Consuming oils with high polyunsaturated fatty acid content, in particular those containing omega-3s, is beneficial for the health. But the mechanisms underlying this phenomenon are poorly known. Researchers at the Institut de Pharmacologie Mol  culaire et Cellulaire (CNRS/Universit   Nice Sophia Antipolis), the Unit   Compartimentation et Dynamique Cellulaires (CNRS/Institut Curie/UPMC), the INSERM and the Universit   de Poitiers investigated the effect of lipids bearing polyunsaturated chains when they are integrated into cell membranes. Their work shows that the presence of these lipids makes the membranes more malleable and therefore more sensitive to deformation and fission by proteins. These results, published on August 8, 2014 in *Science*, could help explain the extraordinary efficacy of endocytosis in neuron cells.

Consuming polyunsaturated [fatty acids](#) (such as [omega-3 fatty acids](#)) is good for the health. The effects range from [neuronal differentiation](#) to protection against cerebral ischemia. However the molecular mechanisms underlying these effects are poorly understood, prompting researchers to focus on the role of these fatty acids in [cell membrane](#) function.

For a cell to function properly, the membrane must be able to deform and divide into small vesicles. This phenomenon is called endocytosis. Generally, these vesicles allow the cells to encapsulate molecules and transport them.. In neurons, these [synaptic vesicles](#) will act as a transmission pathway to the synapse for nerve messages. They are formed inside the cell, then they move to its exterior and fuse with its

membrane, to transmit the neurotransmitters that they contain. Then they reform in less than a tenth of a second: this is synaptic recycling.

In the work published in *Science*, the researchers show that cell- or artificial membranes rich in polyunsaturated lipids are much more sensitive to the action of two proteins, dynamin and endophilin, which facilitate membrane deformation and fission. Other measurements in the study and in simulations suggest that these lipids also make the membranes more malleable. By facilitating the deformation and scission necessary for endocytosis, the presence of polyunsaturated lipids could explain rapid synaptic vesicle recycling.. The abundance of these lipids in the brain could then represent a major advantage for cognitive function.

This work partially sheds light on the mode of action of omega-3. Considering that the body cannot synthesize them and that they can only be supplied by a suitable diet (rich in oily fish, etc.), it seems important to continue this work to understand the link between the functions performed by these lipids in the neuronal membrane and their health benefits.

More information: M. Pinot, et al. Polyunsaturated phospholipids facilitate membrane deformation and fission by endocytic proteins. *Science*, 2014; 345 (6197): 693 [DOI: 10.1126/science.1255288](https://doi.org/10.1126/science.1255288)

Provided by CNRS

Citation: Lipids boost the brain (2014, August 11) retrieved 5 May 2024 from <https://medicalxpress.com/news/2014-08-lipids-boost-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.