

Past pain patterns can help predict future low back pain

August 10 2014




Image courtesy of Blausen Medical

(HealthDay)—Repeated measurements of low back pain (LBP)-related variables are necessary to identify patterns in a fluctuating condition, according to a study published in the Aug. 1 issue of *The Spine Journal*.

Iben Axén, Ph.D., from the Karolinska Institutet in Stockholm, and colleagues collected the "number of days with bothersome pain" via weekly text messages for six months from 244 subjects with nonspecific LBP.

The researchers found that half of the subjects (mean age, 44 years) had experienced LBP for more than 30 days in the previous year. The risk of experiencing a day with bothersome LBP varied over time. The previous duration of [pain intensity](#), the presence of [leg pain](#), and the duration of LBP the previous year showed a predictive ability for all time points;

however, pain intensity, leg pain, and self-rated health showed inconsistent predictive patterns.

"These results may explain the diversity of the results of the predictor studies in the literature," the authors write. "Findings from this study indicate that outcomes should be measured much more frequently in order to obtain more accurate appraisal of temporal pain patterns and estimates of [pain](#) predictors."

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2014 [HealthDay](#). All rights reserved.

Citation: Past pain patterns can help predict future low back pain (2014, August 10) retrieved 25 May 2024 from <https://medicalxpress.com/news/2014-08-pain-patterns-future.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
