

Researchers publish first study of brain activation in MS using fNIRS

August 27 2014



Dr. Wylie of Kessler Foundation demonstrates the portable device for recording functional near infrared spectroscopy in cognitive research studies of brain activation (NIRScout, NIRx Medical Technologies, LLC). Credit: Kessler Foundation

Using functional near infrared spectroscopy (fNIRS), Kessler Foundation researchers have shown differential brain activation patterns

between people with multiple sclerosis (MS) and healthy controls. This is the first MS study in which brain activation was studied using fNIRS while participants performed a cognitive task.

The article, "Neuroimaging and cognition using functional near [infrared spectroscopy](#) (fNIRS) in [multiple sclerosis](#)," was published online on June 11 by *Brain Imaging and Behavior*. Authors are Jelena Stojanovic-Radic, PhD, Glenn Wylie, DPhil, Gerald Voelbel, PhD, Nancy Chiaravalloti, PhD, and John DeLuca, PhD.

Researchers compared 13 individuals with MS with 12 controls for their performance on a working memory task with four levels of difficulty. Most such studies have employed functional magnetic resonance imaging (fMRI); fNIRS has been used infrequently in clinical populations, and has not been applied previously to neuroimaging research in MS. Studies comparing fMRI findings with those of fNIRS, however, show broad agreement in terms of activation patterns.

Results showed differences in activation between the groups that were dependent on task load. The MS group had an increase in activation at low [task difficulty](#) and a decrease in activation at high task difficulty. Conversely, in the control group, activation decreased with low task difficulty and increased with high task difficulty. Performance accuracy was lower in the MS group for low task load; there were no differences between the groups at the higher task loads.

"The data we obtained via fNIRS are consistent with fMRI data for clinical populations. We demonstrated that fNIRS is capable of detecting neuronal activation with a reasonable degree of detail," noted Glenn Wylie, DPhil, associate director of Neuroscience Research and the Neuroimaging Center at Kessler Foundation. "We attribute the differences in [brain activation patterns](#) to the effort expended during the working memory task rather than to differences in speed of processing,"

he added. "Because fNIRS is more portable and easier to use than fMRI, it may offer advantages in monitoring cognitive interventions that require frequent scans."

In addition to working memory, future research in clinical populations should focus on processing speed and episodic memory, cognitive functions that are also affected in MS.

Provided by Kessler Foundation

Citation: Researchers publish first study of brain activation in MS using fNIRS (2014, August 27) retrieved 20 March 2024 from <https://medicalxpress.com/news/2014-08-publish-brain-ms-fnirs.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--