

# Social inequalities in salt consumption remain

August 26 2014

---



People from low socio-economic positions in Britain still eat more salt than those from higher socio-economic positions, irrespective of where they live.

A paper published in the *BMJ Open* journal and led by Warwick Medical School suggests social inequalities in salt intake have hardly changed in the period from 2000-01 to 2011. This is despite a national average salt reduction over this time.

This paper is the first to monitor social inequalities following the national salt reduction programme.

The research was carried out by the World Health Organization Collaborating Centre for Nutrition, based at Warwick Medical School, University of Warwick.

Researchers looked at the geographical distribution of habitual dietary salt intake in Britain and its association with manual occupations and educational attainments, both indicators of socio-economic position and key determinants of health.

The researchers used the British National Diet and Nutrition Survey (2008-11), a national representative sample of 1,027 men and women aged 19-64 years living in Britain. Salt intake was assessed with a 4-day food diary, assessing [salt consumption](#) from food. Salt added by consumers at the table and during cooking (discretionary salt) was not measured.

There was a significant reduction in dietary salt consumption from 2000-1 to 2008-11 of 0.9g of salt per day, consistent with the total reduction in salt consumption of 1.4g per day reported nationally when also discretionary use of salt is taken into account.

Professor Francesco Cappuccio, senior author and Director of the WHO Collaborating Centre, said: "Whilst we are pleased to record an average national reduction in salt consumption coming from food of nearly a gram per day, we are disappointed to find out that the benefits of such a programme have not reached those most in need. These results are important as people of low socio-economic background are more likely to develop high blood pressure (hypertension) and to suffer disproportionately from strokes, heart attacks and renal failure."

Professor Cappuccio continued: "The diet of disadvantaged socio-economic groups tends to be made up of low-quality, salt-dense, high-fat, high-calorie unhealthy cheap foods. We have seen a reduction in salt

intake in Britain thanks to a policy, which included awareness campaigns, food reformulation and monitoring. However, clearly poorer households still have less healthy shopping baskets and the broad reformulation of foods high in [salt](#) has not reached them as much as we would have hoped."

"In our continued effort to reduce population [salt intake](#) towards a 6g per day target in Britain, it is crucial to understand the reasons for these [social inequalities](#) so as to correct this gap for an equitable and cost-effective delivery of cardiovascular prevention", Professor Cappuccio concluded.

Provided by University of Warwick

Citation: Social inequalities in salt consumption remain (2014, August 26) retrieved 25 April 2024 from <https://medicalxpress.com/news/2014-08-social-inequalities-salt-consumption.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.