


New research to help predict teen suicide

August 8 2014



Associate Professor James Scott

Health practitioners will be better equipped to predict self-harm or suicide attempts in adolescents as a result of a University of Queensland study on psychosis in young people.

The UQ Centre for Clinical Research's Associate Professor James Scott said the research showed that adolescents who had psychotic experiences were 13 times more likely to attempt suicide and 11 times more likely to self-harm in the following 12 months than those who had never had a psychotic episode.

"Psychotic experiences and [psychological distress](#) should serve as

warning signs that a young person is at high risk of self-harm or attempting suicide," he said.

"We would encourage health practitioners to ask young people about [psychotic symptoms](#) such as hearing voices or feeling paranoid when they are assessing their mental health and the likelihood of self-harm.

"This would ensure that the appropriate support, health resources and adequate risk-management plans are in place to reduce [suicide attempts](#) in this age group."

The study is believed to be the first of its kind to examine whether psychotic experiences, psychological distress (high levels of anxiety or depression) or both are associated with self-harm and suicide attempts.

"It's also known that approximately one in every 12 adolescents has psychotic experiences, or odd or unusual beliefs such as paranoid ideas or hallucinations," Associate Professor Scott said.

"These symptoms are much more common than was once thought.

"Suicide is the leading cause of death among young people in Australia, and non-suicidal self-harm occurs in approximately one in every seven [young people](#), so it is crucial that this issue is addressed."

The study found that teens who had experienced psychological distress but not [psychotic experiences](#) were three times more likely to [self-harm](#) over the following 12 months and five times more likely to attempt suicide than those who had not experienced psychological distress.

The research involved 1975 Australians aged 12 to 17 over a 12-month period.

Provided by University of Queensland

Citation: New research to help predict teen suicide (2014, August 8) retrieved 2 May 2024 from <https://medicalxpress.com/news/2014-08-teen-suicide.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.