

Risk of esophageal cancer decreases with height

September 25 2014

Taller individuals are less likely to develop esophageal cancer and its precursor, Barrett's esophagus, according to a new study¹ in [Clinical Gastroenterology and Hepatology](#), the official clinical practice journal of the American Gastroenterological Association.

"Individuals in the lowest quartile of [height](#) (under 5'7" for men and 5'2" for women) were roughly twice as likely as individuals in the highest quartile of height (taller than 6' for men and 5'5" for women) to have Barrett's [esophagus](#) or esophageal cancer," said Aaron P. Thrift, PhD, lead study author from the Public Health Sciences Division, Fred Hutchinson Cancer Research Center, Seattle, WA. "Interestingly, the relationship between height and esophageal cancer is opposite from many other cancers—including colorectal, prostate and breast—where greater height is associated with an increased risk."

Researchers conducted a large pooled analysis using data from 14 population-based epidemiologic studies within the International Barrett's and Esophageal Adenocarcinoma Consortium (BEACON), including 1,000 cases of esophageal cancer and twice as many cases of Barrett's esophagus, and twice as many controls. The researchers conducted multiple analyses, including using Mendelian randomization (which incorporates genetic information with traditional approaches) to overcome issues of confounding and bias. The results from all analyses consistently demonstrated an inverse association between height and Barrett's esophagus or esophageal cancer. There were no differences in these estimates based on sex, age, education, smoking, GERD symptoms

or body mass index. Adjusting for abdominal obesity yielded similar results.

"The identification of risk factors, such as height, will allow us to create more sophisticated and accurate methods to quantify patient risk, which will hopefully be used in the future to decide who should undergo endoscopic screening for these conditions," added Dr. Thrift.

The researchers report no obvious explanation for the association between short height and Barrett's esophagus or esophageal cancer. Future studies investigating the potential causal mechanisms by which risk for Barrett's esophagus or [esophageal cancer](#) might be influenced by height are justified.

Esophageal cancer incidence increased eight-fold in the U.S. from 1973 to 2008. Almost all cases arise from Barrett's esophagus. Learn more about the management of Barrett's esophagus in the [American Gastroenterological Association medical position statement](#).

More information: Thrift, A. P., Risk of Esophageal Adenocarcinoma Decreases With Height, Based on Consortium Analysis and Confirmed by Mendelian Randomization. *Clinical Gastroenterology and Hepatology* 2014; 12(10): 1667-1676.e1

Provided by American Gastroenterological Association

Citation: Risk of esophageal cancer decreases with height (2014, September 25) retrieved 26 April 2024 from <https://medicalxpress.com/news/2014-09-esophageal-cancer-decreases-height.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.