

A 'frenemy' in Parkinson's disease takes to crowdsourcing

September 29 2014

The protein alpha-synuclein is a well-known player in Parkinson's disease and other related neurological conditions, such as dementia with Lewy bodies. Its normal functions, however, have long remained unknown. An enticing mystery, say researchers, who contend that understanding the normal is critical in resolving the abnormal.

Alpha-synuclein typically resides at presynaptic terminals – the communication hubs of neurons where neurotransmitters are released to other neurons. In previous studies, Subhojit Roy, MD, PhD, and colleagues at the University of California, San Diego School of Medicine had reported that <u>alpha-synuclein</u> diminishes neurotransmitter release, suppressing communication among neurons. The findings suggested that alpha-synuclein might be a kind of singular brake, helping to prevent unrestricted firing by neurons. Precisely how, though, was a mystery.

Then Harvard University researchers reported in a recent study that alpha-synuclein self-assembles multiple copies of itself inside neurons, upending an earlier notion that the protein worked alone. And in a new paper, published this month in *Current Biology*, Roy, a cell biologist and neuropathologist in the departments of Pathology and Neurosciences, and co-authors put two and two together, explaining how these aggregates of alpha-synuclein, known as multimers, might actually function normally inside neurons.

First, they confirmed that alpha-synuclein multimers do in fact congregate at synapses, where they help cluster synaptic vesicles and



restrict their mobility. Synaptic vesicles are essentially tiny packages created by <u>neurons</u> and filled with neurotransmitters to be released. By clustering these vesicles at the synapse, alpha-synuclein fundamentally restricts neurotransmission. The effect is not unlike a traffic light – slowing traffic down by bunching cars at street corners to regulate the overall flow.

"In normal doses, alpha-synuclein is not a mechanism to impair communication, but rather to manage it. However it's quite possible that in disease, abnormal elevations of alpha-synuclein levels lead to a heightened suppression of neurotransmission and synaptic toxicity," said Roy.

"Though this is obviously not the only event contributing to overall disease neuropathology, it might be one of the very first triggers, nudging the synapse to a point of no return. As such, it may be a neuronal event of critical therapeutic relevance."

Indeed, Roy noted that alpha-synuclein has become a major target for potential drug therapies attempting to reduce or modify its levels and activity.

Provided by University of California - San Diego

Citation: A 'frenemy' in Parkinson's disease takes to crowdsourcing (2014, September 29) retrieved 30 April 2024 from https://medicalxpress.com/news/2014-09-frenemy-parkinson-disease-crowdsourcing.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.