

# Medications plus parent training may help kids with aggression, ADHD

September 12 2014, by Maureen Salamon, Healthday Reporter

---



Combination treatment seems to reduce anger and violent tendencies, study finds.

(HealthDay)—Combining two medications with parent training appears to improve anger, irritability and violent tendencies in children whose attention-deficit/hyperactivity disorder (ADHD) is coupled with severe aggression, a new study suggests.

"Augmented" therapy that consists of stimulant and antipsychotic drugs, along with parent training in [behavioral management](#) techniques, was rated more effective by parents than "basic" therapy pairing only the stimulant and [parent training](#), researchers found.

"An important finding of this study was that at the end of nine weeks, approximately half of all [children](#) receiving basic therapy were still rated

by their parents as being impaired... with symptoms interfering with school or social functioning," said study author Kenneth Gadow, a professor of psychiatry at Stony Brook University in New York.

"In the augmented group receiving three interventions for aggression, about one-quarter still, at the end of nine weeks, were rated by their parents as being impaired," he added. "And that suggests, even with highly effective therapies, that many of these children still have unmet treatment needs."

The drugs used in the study, published in the September issue of the *Journal of the American Academy of Child and Adolescent Psychiatry*, included the widely prescribed ADHD stimulant Concerta (methylphenidate) and the antipsychotic Risperdal (risperidone).

Approximately 11 percent of American children aged 4 to 17 have been diagnosed with ADHD, which includes symptoms such as impulsivity, hyperactivity, and difficulty focusing and controlling behavior, according to the U.S. Centers for Disease Control and Prevention.

As many as half of children with ADHD also display significant, disruptive aggression, according to an editorial accompanying the new research.

"This is very common among kids with ADHD, but unfortunately it complicates the picture for treatment," said Erin Schoenfelder, a clinical psychologist at Seattle Children's Hospital and assistant professor of psychiatry and behavioral health at University of Washington. "It really is staggeringly high."

In the new research, Gadow and his colleagues divided a group of nearly 170 children aged 6 to 12 with ADHD and aggression problems into two treatment groups—basic and augmented. The basic group received

Concerta and their parents underwent behavioral management training.

The augmented group received Concerta and parental training as well, but also took the antipsychotic Risperdal. Both groups were followed for nine weeks.

While both groups of children displayed marked reduction in symptoms, improvement ratings varied depending on whether parents or teachers were making the assessment. Parents reported that children on augmented therapy were less likely to be impaired socially or academically by their anger and irritability than children on basic therapy. On the other hand, teachers found few differences in these measures.

Instead, teachers of those on augmented therapy reported significant drops in ADHD symptoms, especially impulsivity, compared with teachers of children on basic therapy.

Gadow and Schoenfelder agreed that the conflicting parent-teacher ratings demonstrate a familiar concept: that children's behaviors vary in different settings, whether or not they have ADHD.

"Just like adults, they adapt their behaviors to be more appropriate for the setting they're in," Gadow said. "People do differ, however, in their ability to modify their behavior from one setting to the next, and some children are much more variable [in this regard]."

Schoenfelder said long-term evidence is needed indicating that combining Concerta and Risperdal is safe in children, but "it appears from this study that the combination was well-tolerated and something practitioners could [adjust the dosage of] effectively."

"Folks trying this type of treatment should have close monitoring," she

added. "This is a starting point. It's a combination doctors may try when they see this blend of aggressive and hyperactive behaviors. It certainly will require adjustment . . . but it's exciting to find something helpful for a significant proportion of the kids studied."

The study authors pointed out that their findings only apply to children with ADHD who exhibit severe irritability and aggression. They noted that the study's findings are not generalizable for the ADHD population as a whole.

**More information:** The U.S. Centers for Disease Control and Prevention offers more information about [ADHD](#).

Copyright © 2014 [HealthDay](#). All rights reserved.

Citation: Medications plus parent training may help kids with aggression, ADHD (2014, September 12) retrieved 4 May 2024 from <https://medicalxpress.com/news/2014-09-medications-parent-kids-aggression-adhd.html>

|  |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|