

# Phthalates heighten risk for childhood asthma

September 17 2014

---

Researchers at the Columbia Center for Children's Environmental Health at the Mailman School of Public Health are the first to demonstrate an association between childhood asthma and prenatal exposure to two phthalates used in a diverse array of household products. Results appear online in the journal *Environmental Health Perspectives*.

Children born to [mothers](#) exposed during pregnancy to higher levels of the chemicals, butylbenzyl phthalate (BBzP) and di-n-butyl phthalate (DnBP) had a 72 percent and 78 percent increase in risk of developing asthma between age 5 and 11, respectively, compared with children of mothers with lower levels of exposure, the researchers found.

"Everyone from parents to policymakers is concerned by the steep rise in the number of children who develop asthma. Our goal is to try and uncover causes of this epidemic so we can better protect young children from this debilitating condition," says first author Robin Whyatt, DrPH, professor of Environmental Health Sciences and co-deputy director of the Columbia Center for Children's Environmental Health at the Mailman School.

"Our study presents evidence that these two phthalates are among a range of known risk factors for asthma," adds Dr. Whyatt. Other risk factors include tobacco smoke, air pollution, obesity, and a history of allergies.

Phthalates are used in everything from synthetic fragrances to plastic

food containers, vinyl flooring, insect repellent, shower curtains, even steering wheels and dashboards ("new car smell" contains phthalates). Since 2009, several phthalates—including BBzP and DnBP—have been banned from children's toys and other childcare articles in the United States. However, no steps have been taken to protect the developing fetus by alerting pregnant women to potential exposures. In the U.S., phthalates are rarely listed as ingredients on products in which they are used.

"The fetus is extremely vulnerable during pregnancy. While it is incumbent on mothers to do everything they can to protect their child, they are virtually helpless when it comes to phthalates like BBzP and DnBP that are unavoidable. If we want to protect children, we have to protect pregnant women," says the study's senior author Rachel Miller, MD, professor of Medicine (in Pediatrics) and Environmental Health Sciences; and chief of Pediatric Allergy, Immunology, and Rheumatology at Columbia University Medical Center; and co-deputy director of the Columbia Center for Children's Environmental Health at Columbia's Mailman School of Public Health.

The researchers followed 300 [pregnant women](#) and their children enrolled in the Center's longitudinal cohort study in New York City. Levels of exposure to four suspect phthalates were measured through chemical metabolites in urine. Samples were collected from mothers during the third trimester and in children at ages 3, 5, and 7. Chemical analysis was done at the Centers for Disease Control and Prevention. The researchers compared asthma risk in the children between ages 5 and 11 year who were born to mothers with concentrations of the phthalates in the upper third versus the lower third.

Almost a third of the children (94) developed physician-diagnosed asthma. Another 60 children had a history of wheeze and other asthma-like symptoms without a diagnosis of asthma. In this group too, the

researchers found a link between [prenatal exposure](#) to DnBP and the symptoms. Two phthalates studied—di-2-ethylhexyl phthalate (DEHP) and diethyl phthalate (DEP)—were not associated with a diagnosis of asthma or asthma-like symptoms.

All but one of the urinary samples from mothers and children contained metabolites for all four phthalates. Samples had a wide range of concentrations—from not detected to 550 ng/ml for the BBzP metabolites and from 1 to 1,110 ng/mL for DnBP metabolites. It is unclear what behaviors or products contributed to the greater exposures.

The current findings build on the team's 2012 studies on the same cohort of mothers and children and reporting that [children](#) exposed to DEP or BBzP had elevated risk of asthma-related airway inflammation and prenatal exposure to BBzP was associated with elevated risk of childhood eczema. In 2013, the researchers reported that early childhood exposure to bisphenol A (BPA) also was associated with risk for asthma. More recently, they found that the risk from BPA exposure during childhood increased substantially if the mothers also had higher exposure to BBzP during pregnancy, indicating these endocrine disruptors interact to increase [childhood asthma](#) risk (results were published in August in the *Journal of Allergy and Clinical Immunology*).

For now, it's not known how [phthalates](#) increase risk for [asthma](#). Studies of prenatal exposure to BBzP hint that inflammation and oxidative stress may play a role.

Provided by Columbia University's Mailman School of Public Health

Citation: Phthalates heighten risk for childhood asthma (2014, September 17) retrieved 23 April 2024 from <https://medicalxpress.com/news/2014-09-phthalates-heighten-childhood-asthma.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.