

Review: Rapid antigen tests accurate for strep diagnosis

September 11 2014



Image courtesy of Blausen Medical

(HealthDay)—Rapid antigen diagnostic tests (RADTs) can be used for accurate diagnosis of group A streptococcal (GAS) pharyngitis for management of sore throat in primary care settings, according to a study published online Sept. 8 in *Pediatrics*.

Wei Ling Lean, M.B.B.S., from Royal Children's Hospital in Australia, and colleagues conducted a systematic literature review to identify studies evaluating the [diagnostic accuracy](#) of RADTs for GAS pharyngitis.

The researchers identified 48 studies that assessed the diagnostic accuracy of GAS RADTs using throat culture on blood agar as a reference standard. Sensitivity and specificity of RADTs using overall summary estimates were 0.86 and 0.96, respectively. Estimates for

pediatric data were similar. The best diagnostic accuracy was seen with molecular-based RADTs. Study methodologies varied considerably. Insufficient data prevented meta-regression/subgroup analysis within each test type.

"RADTs may not require culture backup for negative tests in most low-incidence rheumatic fever settings," the authors write.

Two study authors received funding for a 2012 study of a Quidel Corp. rapid antigen diagnostic test product.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2014 [HealthDay](#). All rights reserved.

Citation: Review: Rapid antigen tests accurate for strep diagnosis (2014, September 11) retrieved 23 April 2024 from
<https://medicalxpress.com/news/2014-09-rapid-antigen-accurate-strep-diagnosis.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--