

Monitoring the response of bone metastases to treatment using MRI and PET

September 10 2014

Imaging technologies are very useful in evaluating a patient's response to cancer treatment, and this can be done quite effectively for most tumors using RECIST, Response Evaluation Criteria in Solid Tumors. However, RECIST works well for tumors located in soft tissue, but not so well for cancers that spread to the bone, such as is the case for prostate and breast cancers. More effort, therefore, is needed to improve our understanding of how to monitor the response of bone metastases to treatment using magnetic resonance imaging (MRI) and positron emission tomography (PET), and a recent EORTC Imaging Group review and position statement published in the *European Journal of Cancer* is a decidedly welcome contribution.

The paper published by these EORTC Imaging Group investigators highlights the most recent developments in MRI and PET and elucidates how these techniques can be used to detect [bone metastases](#) at an early stage as well as monitor the response of these to treatment. They describe the current state of the art in PET and MRI, the strengths, weaknesses and complementarity of various imaging techniques with respect to specific indications, and recommendations for choosing the most appropriate imaging technique.

Prof. Frederic Lecouvet of the Cliniques Universitaires Saint Luc in Brussels and lead author of this review and position statement says, "Assessing the response of metastases to treatment is something we do on a day to day basis, both in our oncology practice as well as for clinical trials. If the metastases occur in soft tissues, then we already have

validated criteria, such as RECIST, that enable us to evaluate the response. A host of problems, however, limits our ability to measure the response to treatment of metastases found in [bone](#). These range from the characteristics of bone metastatic disease, the structure of bone itself, to the sensitivity, specificity, and resolution of imaging methods available until now. We hope to resolve these issues, and our review is a step in the right direction."

Techniques such as MRI and PET have the potential to detect the spreading of a cancer into the bone marrow at an earlier stage and also determine the extent of this spread, and the efforts of the EORTC researchers will go a long way towards making these techniques a routine part of oncology practice.

Provided by European Organisation for Research and Treatment of Cancer

Citation: Monitoring the response of bone metastases to treatment using MRI and PET (2014, September 10) retrieved 18 April 2024 from <https://medicalxpress.com/news/2014-09-response-bone-metastases-treatment-mri.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--