

Results of CLEAN-TAVI trial reported at TCT 2014

September 15 2014

A first-of-its kind study found that using a cerebral protection device during transcatheter aortic valve replacement (TAVR) can significantly reduce the number and volume of cerebral lesions in high risk patients with severe aortic stenosis.

Findings were reported today at the 26th annual Transcatheter Cardiovascular Therapeutics (TCT) scientific symposium. Sponsored by the Cardiovascular Research Foundation (CRF), TCT is the world's premier educational meeting specializing in interventional cardiovascular medicine.

Stroke remains a major predictor of mortality after TAVR. Cerebral protection devices are designed to reduce the risk of stroke by capturing and removing embolic debris that can be released during the procedure. CLEAN-TAVI was a prospective, randomized, double-blind single center study, and the first trial to examine the impact of a cerebral protection device in preventing MRI-detected brain lesions during TAVR. The primary endpoint was the number of lesions in the protected brain region as determined by diffusion-weighted magnetic resonance imaging (DW-MRI) subtraction at two days post TAVR. Secondary endpoints included the total lesion volume at two and seven days after TAVR and lesion number at seven days.

A total of 100 patients with [severe aortic stenosis](#) who were at increased surgical risk were recruited and randomly assigned in a 1:1 ratio to TAVR with cerebral protection (device group) or TAVR alone ([control](#)

[group](#)). Patients underwent MRIs of the brain before and at two and seven days after TAVR.

After two days, the median number of lesions in the protected regions in the device group was significantly lower than the control group (4 vs. 10, respectively, $p=0.009$). At seven days after TAVR, the median lesion number was also significantly lower in the device group compared to the control group (3 vs. 7, $p=0.0023$). In addition, the median total lesion volume in the protected area was significantly smaller in the device group compared to the control group at two days (246 mm³ vs. 527 mm³, respectively, $p=0.0023$) and at seven days (101 mm³ vs. 292 mm³, $p=0.002$).

"In patients with severe aortic stenosis who are at increased surgical risk, the use of a cerebral protection device during TAVR significantly reduced the number of cerebral lesions in the protected brain regions," said lead investigator Axel Linke, MD. Dr Linke is a Professor at the University of Leipzig Heart Center in Germany.

"Device use also reduced the volume of cerebral lesions as determined by DW-MRI."

More information: The results of the CLEAN-TAVI trial was presented on Saturday, September 13 at 11:42 AM EDT in the Main Arena (Level 3, Ballroom) of the Walter E. Washington Convention Center.

Provided by Cardiovascular Research Foundation

Citation: Results of CLEAN-TAVI trial reported at TCT 2014 (2014, September 15) retrieved 5 May 2024 from <https://medicalxpress.com/news/2014-09-results-clean-tavi-trial-tct.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.