

# Report reveals dangerous gap in understanding of hepatitis C transmission

September 18 2014, by Fran Strachan

---



Credit: Thinkstock

A wide-ranging report by UNSW's Centre for Social Research in Health shows hepatitis C knowledge in young people and gay men is dangerously low.

Findings from the Annual Report of Trends in Behaviour (ARTB 2014) show that NSW young people who use drugs, or are exposed to injecting drugs, do not have enough knowledge of hepatitis C to protect themselves against the disease.

The results of the online survey of 757 young people across NSW aged

16–26 show that 40% of respondents reported illicit drug use in the previous 12 months and a small proportion of respondents reported use of drugs that could be injected, such as cocaine (8%), methamphetamine (6%), heroin (0.4%), and other opioids (1%).

Eleven percent of respondents reported exposure to injecting in the previous 12 months, either through close friends or a partner who injected, or having been offered drugs to inject.

Centre for Social Research in Health Deputy Director, Professor Carla Treloar, said the association of hepatitis C with [injecting drug](#) use makes it difficult to deliver information to young people in school settings or via general media.

"Our research with older people who use drugs tells us that they often don't find out about hepatitis C until some time after they have started to inject drugs," said Professor Treloar.

Young people who have recently commenced injecting drugs, or spend time with others who use drugs are a priority population for hepatitis C prevention, because of the high risk of acquiring hepatitis C shortly after initiating injecting.

"This research tells us that we need find innovative ways to connect [young people](#) with information that is credible and relevant to them," Professor Treloar said.

## **Gay men – Injecting drug use and hepatitis C knowledge**

A 2013 Centre for Social Research in Health national, [online survey](#) of 474 gay and bisexual men on injecting drug use and hepatitis C testing

shows that 15% of men injected drugs in the six months prior to the survey .

Crystal methamphetamine was the drug most commonly injected (75%) and was typically injected before or during sex (95%).

A high proportion of men reported reusing and passing on needles, syringes and other injecting equipment in the six months prior to the study (41%), and sharing commonly occurred with regular and casual sex partners. Sharing of injecting equipment remains the primary risk factor for hepatitis C transmission.

The survey also shows a high percentage (80%) of gay men have tested for hepatitis C and 58% reported testing in the previous 12 months. However, barriers to testing included having never been offered a test, low risk perception and uncertainties about practical aspects, such as what a test involved and where to obtain one.

"These findings suggest a need for continued dissemination of harm reduction information in relation to the risk of acquiring [hepatitis C](#), as well as a need for further research into gay men's injecting practices to guide harm reduction responses," said Centre for Social Research in Health Director John de Wit.

"However, this research, combined with the findings from our Gay Community Periodic Survey, show that there is a slow but ongoing decline in injecting [drug](#) use in [gay men](#)."

Provided by University of New South Wales

Citation: Report reveals dangerous gap in understanding of hepatitis C transmission (2014, September 18) retrieved 27 April 2024 from <https://medicalxpress.com/news/2014-09-reveals->

[dangerous-gap-hepatitis-transmission.html](https://www.medicalxpress.com/dangerous-gap-hepatitis-transmission.html)

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.