

Ultrasound can accurately diagnose carpal tunnel

September 30 2014



Image courtesy of Blausen Medical

(HealthDay)—Ultrasound can accurately confirm the diagnosis of carpal tunnel syndrome, according to a study published in the Sept. 3 issue of *The Journal of Bone & Joint Surgery*.

John R. Fowler, M.D., from the University of Pittsburgh, and colleagues compared the sensitivity and specificity of [ultrasound](#) and electrodiagnostic testing for diagnosing [carpal tunnel syndrome](#) in a series of 85 patients. All patients were assessed with the Carpal Tunnel Syndrome 6 (CTS-6) clinical diagnostic tool, which was used as the reference standard.

The researchers found that ultrasound had a sensitivity of 89 percent and specificity of 90 percent, while electrodiagnostic testing had a sensitivity of 89 percent and specificity of 80 percent. The positive predictive

values were 94 and 89 percent, respectively, for ultrasound and electrodiagnostic testing, while the negative predictive values were 82 and 80 percent, respectively. Ultrasound was accurate in 89 percent of cases and electrodiagnostic testing was accurate in 86 percent of cases.

"While ultrasound will not replace electrodiagnostic testing in complicated or unclear cases, in a select group of patients with a positive CTS-6, ultrasound can be used to confirm the [diagnosis](#) of carpal tunnel syndrome with better specificity and equal sensitivity as compared with those of electrodiagnostic testing," the authors write.

More information: [Full Text](#)

Copyright © 2014 [HealthDay](#). All rights reserved.

Citation: Ultrasound can accurately diagnose carpal tunnel (2014, September 30) retrieved 26 April 2024 from

<https://medicalxpress.com/news/2014-09-ultrasound-accurately-carpal-tunnel.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.