

White matter measure predicts longer concussion recovery

September 11 2014



(HealthDay)—A measure of white matter in the brain, particularly in males, is an independent predictor of longer time to symptom resolution (TSR) after mild traumatic brain injury (mTBI), according to a study published in the September issue of *Radiology*.

Saeed Fakhran, M.D., from the University of Pittsburgh, and colleagues assessed diffusion-tensor imaging in 69 patients with mTBI (47 male and 22 <u>female patients</u>) and 21 control subjects (10 males and 11 females) whose conventional <u>magnetic resonance images</u> were normal. Fractional anisotropy maps were used as a measure of <u>white matter</u> integrity. Serial neurocognitive testing with Immediate Post-Concussion Assessment and Cognitive Testing was performed in mTBI patients.

The researchers found that compared with female patients with mTBI



and control subjects, male patients with mTBI had significantly lowered fractional anisotropy values in the uncinate fasciculus bilaterally (P 3-month TSR (adjusted odds ratios [OR], 2.27 and 2.38; P = 0.04 and P

"Relative sparing of the uncinate fasciculus is seen in female compared with male patients after mTBI, with sex and uncinate fasciculus fractional anisotropy values as stronger predictors of TSR than initial symptom severity," the authors write.

More information: Abstract

Full Text (subscription or payment may be required)

Copyright © 2014 HealthDay. All rights reserved.

Citation: White matter measure predicts longer concussion recovery (2014, September 11) retrieved 16 June 2024 from <u>https://medicalxpress.com/news/2014-09-white-longer-concussion-recovery.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.