

# Progression of age-related macular degeneration in one eye then fellow eye

October 23 2014

---

Having age-related macular degeneration (AMD) in one eye was associated with an increased incidence of AMD and accelerated progression of the debilitating disease in the other eye, writes author Ronald E. Gangnon, Ph.D., of the University of Wisconsin School of Medicine and Public Health, Madison, and colleagues.

AMD is thought to be a symmetric disease, although one eye may precede the other in progression.

The authors examined the effect of severity of AMD in one eye on the incidence, progression and regression in the other eye. Data from 4,379 participants in the Beaver Dam Eye Study were used. Retinal photographs were used to assess the incidence, progression and regression of AMD.

While more severe AMD in one eye was associated with increased incidence and accelerated progression in the other eye, less severe AMD in one eye was associated with less progression in the other eye.

"In a cohort that was observed for 20 years, we showed that AMD severity in one eye largely tracks AMD severity in the fellow eye at all stages of the disease. ... Our model demonstrated the effect of one eye on the incidence and progression of AMD in its fellow [eye](#) across the entire continuum of AMD severity."

**More information:** *JAMA Ophthalmol.* Published online October 23,

2014. [DOI: 10.1001/jamaophthalmol.2014.4252](https://doi.org/10.1001/jamaophthalmol.2014.4252)

Provided by The JAMA Network Journals

Citation: Progression of age-related macular degeneration in one eye then fellow eye (2014, October 23) retrieved 4 May 2024 from <https://medicalxpress.com/news/2014-10-age-related-macular-degeneration-eye-fellow.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.