

Blood tests predict kidney disease patients' risk of developing heart failure

October 2 2014

Two blood markers are strongly linked with the development of heart failure in individuals with mild to severe kidney disease, according to a study appearing in an upcoming issue of the *Journal of the American Society of Nephrology (JASN)*. Elevations in these markers may indicate subclinical cardiovascular changes that subsequently contribute to the development of heart failure.

Patients with chronic [kidney disease](#) (CKD) are at increased risk of developing heart failure and other cardiovascular diseases. Nisha Bansal, MD, MAS (University of Washington) Amanda Anderson, PhD, MPH (University of Pennsylvania), and their colleagues conducted a study to see if certain blood tests might help identify patients at especially high risk. These tests—which measure proteins called high-sensitivity troponin T (hsTnT) and N-terminal pro-B-type natriuretic peptide (NT-proBNP)—strongly predict heart failure in the general population, but their predictive utility in patients with CKD is unknown. The researchers studied 3483 patients with CKD in the Chronic

Renal Insufficiency Cohort (CRIC) Study who were recruited from June 2003 to August 2008 and were free of heart failure when they enrolled. Patients were followed for a median of nearly 6 years.

Compared with participants with the lowest levels of hsTnT at the start of the study, those with the highest hsTnT levels had a nearly 5-fold higher risk of developing heart failure. Those with the highest NT-proBNP levels had a nearly 10-fold higher risk of developing heart

failure compared with those with the lowest levels.

"This research is important in that it may advance the application of widely available cardiac biomarkers to identify CKD [patients](#) at the highest risk of developing [heart failure](#), the most common cardiovascular complication in this patient population," said Dr. Bansal. "These findings suggest that hsTnT and NT-proBNP may represent distinct biological pathways that likely involve subclinical changes in the structure and function of the heart," said Dr. Anderson.

More information: The article, entitled "High Sensitivity Troponin T and NT-proBNP and Risk of Incident Heart Failure in Patients with Chronic Kidney Disease: The CRIC Study," will appear online at jasn.asnjournals.org/ on October 2, 2014.

Provided by American Society of Nephrology

Citation: Blood tests predict kidney disease patients' risk of developing heart failure (2014, October 2) retrieved 19 April 2024 from <https://medicalxpress.com/news/2014-10-blood-kidney-disease-patients-heart.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--