


A new way to extract bone-making cells from fat tissue

October 6 2014



Finding cells that have bone-making potential is more efficiently done by looking at the genes they express -- ALPL, in this case -- than at proteins on their surface. The bone matrix being produced by cells is stained red in samples of cells that are sorted for not expressing ALPL (left), sorted for expressing ALPL (right), and left as an unsorted control (center). Credit: Darling lab/Brown University

Within our fat lives a variety of cells with the potential to become bone, cartilage, or more fat if properly prompted. This makes adipose tissue, in theory, a readily available reservoir for regenerative therapies such as bone healing if doctors can get enough of those cells and compel them to produce bone.

In a new study in the journal *Stem Cell Research & Therapy*, scientists at Brown University demonstrate a new method for extracting a wide variety of potential bone-producing [cells](#) from human fat. They developed a fluorescent tag that could find and identify cells expressing

a gene called ALPL. Expression of the gene is an indicator of bone-making potential.

If the tag finds the RNA produced when the gene is expressed, it latches on and glows. A machine that detects the fluorescing light then separates out the ALPL-expressing cells.

In the paper, the scientists report that their method produced more than twice the yield of potential bone-makers (9 percent) compared to their best application of another method: sorting cells based on [surface proteins](#) presumed to indicate that a cell is a stem cell (4 percent).

Brown University has applied for a patent on the method of gene expression tagging for producing a tissue.

Meanwhile, the ALPL-expressing cells produced on average more than twice as much bone matrix (and as much as nine times more in some trials) during three weeks of subsequent cultivation than a similar-sized population of unsorted [adipose tissue](#) cells and almost four times more bone matrix than cells that don't express ALPL. ALPL-expressing cells were also better at making [cartilage](#) or fat.

A couple of other research groups have also sorted [stem cells](#) based on gene expression, but they have not done so specifically with the goal of enriching cell populations for a specific tissue, the researchers said.

Lead author and Brown graduate student Hetal Marble said targeting [gene expression](#) rather than surface proteins for the purpose of gathering cells to make a new tissue is a "paradigm shift" in the following regard: Gene expression provides a way to target any cell based on whether it can produce another tissue, while targeting surface proteins limits researchers to harvesting cells that fit a presumed definition of being a stem cell. The new approach, she said, is more pragmatic for the

purpose.

"Approaches like this allow us to isolate all the cells that are capable of doing what we want, whether they fit the archetype of what a stem cell is or not," Marble said. "The paradigm shift is thinking about isolating populations that are able to achieve an end point rather than isolating populations that fit a strictly defined archetype."

In their experiments, though, the team tolerated a four-day delay that they'd like to dispense with in the future. It takes that long for the maximum number of cells to express ALPL when cells are chemically primed to do so.

In future research, said senior author Eric Darling, the Manning Assistant Professor of Molecular Pharmacology, Physiology and Biotechnology and a member of the Center for Biomedical Engineering assistant professor of medical science, the team would like to target a gene expressed much earlier in the differentiation process to see if they can avoid a priming period.

If they can apply the method based on a gene that's expressible within a matter of hours, that could allow future surgeons working on [bone healing](#) to take out some of a patient's fat cells, sort out the best bone-producers (primed or not) and then implant those cells in the bone break within the same surgical session.

"If you can take the patient into the OR, isolate a bunch of their cells, sort them and put them back in that's ideally where we'd like to go with this," Darling said. "Theoretically we could do this with other genes that might upregulate very quickly or are innately expressed.

Provided by Brown University

Citation: A new way to extract bone-making cells from fat tissue (2014, October 6) retrieved 25 April 2024 from <https://medicalxpress.com/news/2014-10-bone-making-cells-fat-tissue.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.