

Healthy lifestyle may cut stroke risk in half for women

October 8 2014



Micrograph showing cortical pseudolaminar necrosis, a finding seen in strokes on medical imaging and at autopsy. H&E-LFB stain. Credit: Nephron/Wikipedia

Women with a healthy diet and lifestyle may be less likely to have a stroke by more than half, according to a study published in the October 8, 2014, online issue of *Neurology*, the medical journal of the American Academy of Neurology.

The study looked at five factors that make up a [healthy lifestyle](#): healthy diet; [moderate alcohol consumption](#); never smoking; physically active; and healthy body mass index (BMI). Compared with [women](#) with none of the five healthy factors, women with all five factors had a 54-percent lower risk of [stroke](#).

"Because the consequences of stroke are usually devastating and irreversible, prevention is of great importance," said study author Susanna C. Larsson, PhD, of the Karolinska Institutet in Stockholm, Sweden. "These results are exciting because they indicate that a healthy diet and lifestyle can substantially reduce the risk of stroke, and these are lifestyle choices that people can make or improve."

For the study, 31,696 Swedish women with an average age of about 60 completed a 350-item questionnaire about their diet and lifestyle. They were then followed for an average of 10 years. A healthy diet was defined as within the top 50 percent of a recommended food score measuring how often the participants ate healthy foods such as fruits, vegetables and low-fat dairy products. Moderate alcohol consumption was defined as three to nine drinks per week. Physically active was defined as walking or biking at least 40 minutes a day along with more vigorous exercise at least one hour per week. Healthy BMI was considered below 25.

Most of the women had two or three of the healthy factors. Only 589 women had all five healthy factors, and 1,535 had none.

There were 1,554 strokes among study participants. The risk of stroke steadily decreased with each additional healthy lifestyle factor.

Women who had a healthier diet were 13 percent less likely to have a type of stroke called a [cerebral infarction](#) than those whose diet was not as healthy. Women with healthier diets had a rate of 28 strokes per

10,000 women per year compared to 43 strokes per 10,000 women per year among those with a less [healthy diet](#).

Cerebral infarction is the most common cause of stroke, accounting for up to 80 to 85 percent of all strokes. Cerebral infarction is caused by a blockage in a blood vessel preventing blood and oxygen from getting to an area of the brain.

There was no relationship between the healthy factors and the risk of [hemorrhagic stroke](#). Hemorrhagic stroke, which is caused by bleeding in and around the brain, accounts for about 15 to 20 percent of all strokes.

Provided by American Academy of Neurology

Citation: Healthy lifestyle may cut stroke risk in half for women (2014, October 8) retrieved 25 April 2024 from <https://medicalxpress.com/news/2014-10-healthy-lifestyle-women.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.