

# Large variation in cesarean rates across US hospitals

October 21 2014

---

Cesarean delivery is the most common inpatient surgery in the United States. US cesarean rates increased from 20.7% in 1996 to 32.9% in 2009 but have since stabilized, with 1.3 million American women having had a cesarean delivery in 2011. Rates of cesarean delivery vary across hospitals, and understanding reasons for the variation could help shed light on practices related to cesarean delivery.

Katy Kozhimannil (University of Minnesota School of Public Health, Minneapolis, USA) and colleagues S.V. Subramanian and Mariana Arcaya (Harvard School of Public Health, Boston, USA) used the 2009 and 2010 US Nationwide Inpatient Sample from the Healthcare Cost and Utilization Project, a 20% sample of US hospitals, to study [hospital](#) variation in [cesarean section](#) rates. They analyzed data for 1,475,457 births in 1,373 hospitals, adjusting rates for maternal diagnoses and sociodemographics, as well as hospital characteristics like size, location, and teaching status.

They found that among all women the cesarean section rate was 33% overall and 22% among women without a prior cesarean section. Rates varied by maternal age, race/ethnicity, and insurance status, and medical diagnoses related to pregnancy and delivery. Variability in hospital cesarean rates did not decrease after adjusting for these maternal factors and hospital characteristics. They found significantly higher variation in hospital rates than would be expected, with the likelihood of cesarean delivery for all women ranging from 19% to 48% across hospitals. Among women without a prior cesarean section, the likelihood of having

a cesarean delivery varied between 11% and 36% across hospitals. They also examined lower risk women, whose likelihood varied between 8% and 32% across hospitals, and higher risk [women](#), whose likelihood varied between 56% and 92% across hospitals. A limitation is that these hospital administrative data, while nationally representative, did not contain information on parity or gestational age, factors that influence cesarean delivery decisions. These measures are collected in other countries such as the United Kingdom but are not routinely included in US hospital discharge summaries. The authors call for these data to be routinely collected and linked to facilitate US research as well as cross-country comparisons.

The authors say, "The findings from this analysis are consistent with studies in Arizona, in Massachusetts, and among births in US military hospitals; however, this analysis is the first to our knowledge to use a nationally representative US sample." They acknowledge the need for future research, "Many other systems- or clinician-based factors may influence use of cesarean section in US hospitals, including liability and insurance factors, the presence and type of a hospital's clinical training program, the role of midwifery, the presence of labor support or birth doulas, individual clinician approaches to labor and delivery management, and practices related to admission and labor management."

**More information:** Kozhimannil KB, Arcaya MC, Subramanian SV (2014) Maternal Clinical Diagnoses and Hospital Variation in the Risk of Cesarean Delivery: Analyses of a National US Hospital Discharge Database. *PLoS Med* 11(10): e1001745. [DOI: 10.1371/journal.pmed.1001745](#)

Provided by Public Library of Science

Citation: Large variation in cesarean rates across US hospitals (2014, October 21) retrieved 24 April 2024 from

<https://medicalxpress.com/news/2014-10-large-variation-cesarean-hospitals.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.