

Low-carbohydrate diet improved quality of life in diabetes patients

October 3 2014

Low-carbohydrate diet has a good effect not only on blood glucose, but also on physical functions, bodily pain and general health, according to a diet study including patients with type 2 diabetes.

The results of a two-year clinical trial in patients with [type 2 diabetes](#), led by Dr Hans Guldbbrand, general practitioner, and Fredrik Nystrom, professor of internal medicine at Linköping University, has previously been published in the journal *Diabetologia*. The trial studied the effects on [blood glucose](#) and blood lipids of a low-carbohydrate [diet](#) compared to a low-fat diet.

The 61 enrolled patients were randomly divided into two groups – one for each diet type and were expected to adhere to the respective diet throughout the study period. It was found that both diet-groups reduced weight equally but the effect on blood glucose was better in the low-carbohydrate group.

The effects of a low-carbohydrate diet and a low-fat diet on wellbeing have now been analysed in a study led by Associate Professor Margareta Bachrach-Lindström. A standardised analysis based on the SF-36 questionnaire was performed. After 12 months in the trial, the low-carbohydrate group improved in regard to the physical component, which includes physical function, bodily pain and [general health](#). No improvements were seen in the low-fat group, despite weight loss. Mental health was similar for both groups and remained unchanged during the study period and did not differ between the groups.

"The result is interesting; it provides an additional argument that a low-carbohydrate diet is beneficial in diabetes," says Hans Guldbrand. "We also found no adverse effects on [mental health](#) with the low-carbohydrate diet, which an earlier study had indicated," he adds.

The interview with the study patients revealed that for both groups there were difficulties adhering to the diet when they ate elsewhere than at home. It could also be problems if not all family members followed the same diet. Both groups expected [health](#) gains by adhering to given dietary advice. The low-carbohydrate diet group expressed that it could be difficult to refrain from potatoes and pasta. The diet for the low-fat group was described as relatively inexpensive and tasty. Benefits of the [low-carbohydrate diet](#) were that the patients felt less hungry and that their appetite for sweets disappeared.

More information: "Randomization to a low-carbohydrate diet advice improves health related quality of life compared with a low-fat diet at similar weight-loss in Type 2 diabetes mellitus," by H. Guldbrand, T. Lindström, B. Dizdar, B. Bunjaku, C.J. Östgren, F.H. Nyström and M. Bachrach-Lindström. *Diabetes Research and Clinical Practice* online 21 September 2014. [DOI: 10.1016/j.diabres.2014.08.032](https://doi.org/10.1016/j.diabres.2014.08.032)

Provided by Linköping University

Citation: Low-carbohydrate diet improved quality of life in diabetes patients (2014, October 3) retrieved 4 May 2024 from <https://medicalxpress.com/news/2014-10-low-carbohydrate-diet-quality-life-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
